

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**

**THIS PAGE BLANK (USPTO,**



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

51) International Patent Classification<sup>6</sup> :

A61K 38/16

A1

(11) International Publication Number:

WO 98/30231

(43) International Publication Date:

16 July 1998 (16.07.98)

(21) International Application Number: PCT/US98/00449

(22) International Filing Date: 7 January 1998 (07.01.98)

## (30) Priority Data:

60/034,905	7 January 1997 (07.01.97)	US
60/055,404	8 August 1997 (08.08.97)	US
60/066,029	14 November 1997 (14.11.97)	US
60/065,442	14 November 1997 (14.11.97)	US

(71) Applicant: AMYLIN PHARMACEUTICALS, INC. [US/US];  
9373 Towne Centre Drive, San Diego, CA 92121 (US).(72) Inventors: BEELEY, Nigel, Robert, Arnold; 227 Loma  
Corta Drive, Solana Beach, CA 92075 (US). PRICKETT,  
Kathryn, S.; 7612 Trailbrush Terrace, San Diego, CA 92126  
(US). BHAVSAR, Sunil; Apartment #7, 917 Torrance  
Street, San Diego, CA 92103 (US).(74) Agents: DUFT, Bradford, J. et al.; Lyon & Lyon LLP, First  
Interstate World Center, Suite 4700, 633 West Fifth Street,  
Los Angeles, CA 90071 (US).(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR,  
BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE,  
GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ,  
LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW,  
MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ,  
TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent  
(GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, ML, MR, NE, SN, TD, TG).

## Published

*With international search report.**Before the expiration of the time limit for amending the  
claims and to be republished in the event of the receipt of  
amendments.*

(54) Title: USE OF EXENDINS AND AGONISTS THEREOF FOR THE REDUCTION OF FOOD INTAKE

## (57) Abstract

Methods for treating conditions or disorders which can be alleviated by reducing food intake are disclosed which comprise administration of an effective amount of an exendin or an exendin agonist, alone or in conjunction with other compounds or compositions that effect satiety. The methods are useful for treating conditions or disorders, including obesity, Type II diabetes, eating disorders, and insulin-resistance syndrome. The methods are also useful for lowering the plasma glucose level, lowering the plasma lipid level, reducing the cardiac risk, reducing the appetite, and reducing the weight of subjects. Pharmaceutical compositions for use in the methods of the invention are also disclosed.

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

USE OF EXENDINS AND AGONISTS THEREOF  
FOR THE REDUCTION OF FOOD INTAKE

5 This application claims the benefit of U.S.  
Provisional Application No. 60/034,905, filed January 7,  
1997, U.S. Provisional Application No. 60/055,404, filed  
August 8, 1997, U.S. Provisional Application No.  
60/066,029 filed November 14, 1997, and U.S. Provisional  
Application No. 60/065,442, November 14, 1997.

10

FIELD OF THE INVENTION

The present invention relates to methods for treating  
conditions or disorders which can be alleviated by  
reducing food intake comprising administration of an  
15 effective amount of an exendin or an exendin agonist alone  
or in conjunction with other compounds or compositions  
that affect satiety such as a leptin or an amylin agonist.  
The methods are useful for treating conditions or  
disorders, in which the reduction of food intake is of  
20 value, including obesity, Type II diabetes, eating  
disorders, and insulin-resistance syndrome. The methods  
are also useful for lowering the plasma lipid level,  
reducing the cardiac risk, reducing the appetite, and  
reducing the weight of subjects. Pharmaceutical  
25 compositions for use in the methods of the invention are  
also disclosed.

BACKGROUND

The following description summarizes information relevant to the present invention. It is not an admission that any of the information provided herein is prior art to the presently claimed invention, nor that any of the publications specifically or implicitly referenced are prior art to that invention.

Exendin

Exendins are peptides that are found in the venom of the Gila-monster, a lizard found in Arizona, and the Mexican Beaded Lizard. Exendin-3 is present in the venom of Heloderma horridum, and exendin-4 is present in the venom of Heloderma suspectum (Eng, J., et al., J. Biol. Chem., 265:20259-62, 1990; Eng, J., et al., J. Biol. Chem., 267:7402-05, 1992). The exendins have some sequence similarity to several members of the glucagon-like peptide family, with the highest homology, 53%, being to GLP-1[7-36]NH<sub>2</sub> (Goke, et al., J. Biol. Chem., 268:19650-55, 1993). GLP-1[7-36]NH<sub>2</sub>, also known as proglucagon[78-107], has an insulinotropic effect, stimulating insulin secretion from pancreatic  $\beta$ -cells; GLP also inhibits glucagon secretion from pancreatic  $\alpha$ -cells (Orskov, et al., Diabetes, 42:658-61, 1993; D'Alessio, et al., J. Clin. Invest., 97:133-38, 1996). GLP-1 is reported to inhibit gastric emptying (Williams B, et al., J. Clin Endocrinol Metab 81 (1): 327-32, 1996; Wettergren A, et al., Dig Dis Sci 38 (4): 665-73, 1993), and gastric acid

secretion. (Schjoldager BT, et al., Diag Dis Sci 34 (5): 703-8, 1989; O'Halloran DJ, et al., J Endocrinol 126 (1): 169-73, 1990; Wettergren A, et al., Diag Dis Sci 38 (4): 665-73, 1993). GLP-1[7-37], which has an additional glycine residue at its carboxy terminus, also stimulates insulin secretion in humans (Orskov, et al., Diabetes, 42:658-61, 1993). A transmembrane G-protein adenylate-cyclase-coupled receptor believed to be responsible for the insulintropic effect of GLP-1 is reported to have been cloned from a  $\beta$ -cell line (Thorens, Proc. Natl. Acad. Sci. USA 89:8641-45 (1992)).

Exendin-4 potently binds at GLP-1 receptors on insulin-secreting  $\beta$ TC1 cells, at dispersed acinar cells from guinea pig pancreas, and at parietal cells from stomach; the peptide is also said to stimulate somatostatin release and inhibit gastrin release in isolated stomachs (Goke, et al., J. Biol. Chem. 268:19650-55, 1993; Schepp, et al., Eur. J. Pharmacol., 69:183-91, 1994; Eissele, et al., Life Sci., 55:629-34, 1994).

Exendin-3 and exendin-4 were reported to stimulate cAMP production in, and amylase release from, pancreatic acinar cells (Malhotra, R., et al., Regulatory Peptides, 41:149-56, 1992; Raufman, et al., J. Biol. Chem. 267:21432-37, 1992; Singh, et al., Regul. Pept. 53:47-59, 1994). The use of exendin-3 and exendin-4 as insulintrophic agents for the treatment of diabetes mellitus and the prevention of hyperglycemia has been proposed (Eng, U.S. Patent No. 5,424,286).

C-terminally truncated exendin peptides such as exendin[9-39], a carboxyamidated molecule, and fragments 3-39 through 9-39 have been reported to be potent and selective antagonists of GLP-1 (Goke, et al., J. Biol. Chem., 268:19650-55, 1993; Raufman, J.P., et al., J. Biol. Chem. 266:2897-902, 1991; Schepp, W., et al., Eur. J. Pharm. 269:183-91, 1994; Montrose-Rafizadeh, et al., Diabetes, 45(Suppl. 2):152A, 1996). Exendin[9-39] is said to block endogenous GLP-1 in vivo, resulting in reduced insulin secretion. Wang, et al., J. Clin. Invest., 95:417-21, 1995; D'Alessio, et al., J. Clin. Invest., 97:133-38, 1996). The receptor apparently responsible for the insulinotropic effect of GLP-1 has reportedly been cloned from rat pancreatic islet cell (Thorens, B., Proc. Natl. Acad. Sci. USA 89:8641-8645, 1992). Exendins and exendin[9-39] are said to bind to the cloned GLP-1 receptor (rat pancreatic  $\beta$ -cell GLP-1 receptor (Fehmann HC, et al., Peptides 15 (3): 453-6, 1994) and human GLP-1 receptor (Thorens B, et al., Diabetes 42 (11): 1678-82, 1993). In cells transfected with the cloned GLP-1 receptor, exendin-4 is reportedly an agonist, i.e., it increases cAMP, while exendin[9-39] is identified as an antagonist, i.e., it blocks the stimulatory actions of exendin-4 and GLP-1. Id.

Exendin[9-39] is also reported to act as an antagonist of the full length exendins, inhibiting stimulation of pancreatic acinar cells by exendin-3 and exendin-4 (Raufman, et al., J. Biol. Chem. 266:2897-902,



1991; Raufman, et al., J. Biol. Chem., 266:21432-37, 1992). It is also reported that exendin[9-39] inhibits the stimulation of plasma insulin levels by exendin-4, and inhibits the somatostatin release-stimulating and gastrin release-inhibiting activities of exendin-4 and GLP-1 (Kolligs, F., et al., Diabetes, 44:16-19, 1995; Eissele, et al., Life Sciences, 55:629-34, 1994).

Exendins have recently been found to inhibit gastric emptying (U.S.S.N. 08/694,954, filed August 8, 1996, which enjoys common ownership with the present invention and is hereby incorporated by reference).

Exendin [9-39] has been used to investigate the physiological relevance of central GLP-1 in control of food intake (Turton, M.D. et al. Nature 379:69-72, 1996).

GLP-1 administered by intracerebroventricular injection inhibits food intake in rats. This satiety-inducing effect of GLP-1 delivered ICV is reported to be inhibited by ICV injection of exendin [9-39] (Turton, supra).

However, it has been reported that GLP-1 does not inhibit food intake in mice when administered by peripheral injection (Turton, M.D., Nature 379:69-72, 1996; Bhavsar, S.P., Soc. Neurosci. Abstr. 21:460 (188.8), 1995).

#### Obesity and Hypernutrition

Obesity, excess adipose tissue, is becoming increasingly prevalent in developed societies. For example, approximately 30% of adults in the U.S. were estimated to be 20 percent above desirable body weight -- an accepted measure of obesity sufficient to impact a health risk (*Harrison's Principles of Internal Medicine*

12th Edition, McGraw Hill, Inc. (1991) p. 411). The pathogenesis of obesity is believed to be multifactorial but the basic problem is that in obese subjects food intake and energy expenditure do not come into balance until there is excess adipose tissue. Attempts to reduce food intake, or hypernutrition, are usually fruitless in the medium term because the weight loss induced by dieting results in both increased appetite and decreased energy expenditure (Leibel et al., (1995) *New England Journal of Medicine* 322: 621-628). The intensity of physical exercise required to expend enough energy to materially lose adipose mass is too great for most people to undertake on a sufficiently frequent basis. Thus, obesity is currently a poorly treatable, chronic, essentially intractable metabolic disorder. Not only is obesity itself believed by some to be undesirable for cosmetic reasons, but obesity also carries serious risk of co-morbidities including, Type 2 diabetes, increased cardiac risk, hypertension, atherosclerosis, degenerative arthritis, and increased incidence of complications of surgery involving general anesthesia. Obesity due to hypernutrition is also a risk factor for the group of conditions called insulin resistance syndrome, or "syndrome X." In syndrome X, it has been reported that there is a linkage between insulin resistance and hypertension. (Watson N. and Sandler M., *Curr. Med. Res. Opin.*, 12(6):374-378 (1991); Kodama J. et al., *Diabetes Care*, 13(11):1109-1111 (1990); Lithell et al., *J. Cardiovasc. Pharmacol.*, 15 Suppl. 5:S46-S52 (1990)).

In those few subjects who do succeed in losing weight, by about 10 percent of body weight, there can be striking improvements in co-morbid conditions, most especially Type 2 diabetes in which dieting and weight loss are the primary therapeutic modality, albeit relatively ineffective in many patients for the reasons stated above. Reducing food intake in obese subjects would decrease the plasma glucose level, the plasma lipid level, and the cardiac risk in these subjects.

Hypernutrition is also the result of, and the psychological cause of, many eating disorders. Reducing food intake would also be beneficial in the treatment of such disorders.

Thus, it can be appreciated that an effective means to reduce food intake is a major challenge and a superior method of treatment would be of great utility. Such a method, and compounds and compositions which are useful therefor, have been invented and are described and claimed herein.

#### SUMMARY OF THE INVENTION

The present invention concerns the surprising discovery that exendins and exendin agonists have a profound and prolonged effect on inhibiting food intake.

The present invention is directed to novel methods for treating conditions or disorders associated with hypernutrition, comprising the administration of an exendin, for example, exendin-3 [SEQ ID NO. 1: His Ser Asp

Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu  
Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro  
Ser Ser Gly Ala Pro Pro Pro Ser], or exendin-4 [SEQ ID NO.  
2: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser], or other  
compounds which effectively bind to the receptor at which  
exendin exerts its action on reducing food intake. These  
methods will be useful in the treatment of, for example,  
10 obesity, diabetes, including Type II or non-insulin  
dependent diabetes, eating disorders, and insulin-  
resistance syndrome.

In a first aspect, the invention features a method of  
treating conditions or disorders which can be alleviated  
15 by reducing food intake in a subject comprising  
administering to the subject a therapeutically effective  
amount of an exendin or an exendin agonist. By an "exendin  
agonist" is meant a compound that mimics the effects of  
exendin on the reduction of food intake by binding to the  
20 receptor or receptors where exendin causes this effect.  
Preferred exendin agonist compounds include those described  
in United States Provisional Patent Application Serial No.  
60/055,404, entitled, "Novel Exendin Agonist Compounds,"  
filed August 8, 1997; United States Provisional Patent  
25 Application Serial No. 60/065,442, entitled, "Novel Exendin  
Agonist Compounds," filed November 14, 1997; and United  
States Provisional Patent Application Serial No.  
60/066,029, entitled, "Novel Exendin Agonist Compounds,"  
filed November 14, 1997; all of which enjoy common

ownership with the present application and all of which are incorporated by this reference into the present application as though fully set forth herein. By "condition or disorder which can be alleviated by reducing food intake" is meant any condition or disorder in a subject that is either caused by, complicated by, or aggravated by a relatively high food intake, or that can be alleviated by reducing food intake. Such conditions or disorders include, but are not limited to, obesity, diabetes, including Type II diabetes, eating disorders, and insulin-resistance syndrome.

Thus, in a first embodiment, the present invention provides a method for treating conditions or disorders which can be alleviated by reducing food intake in a subject comprising administering to said subject a therapeutically effective amount of an exendin or an exendin agonist. Preferred exendin agonist compounds include those described in U.S. Provisional Patent Application Serial Nos. 60/055,404; 60/065,442; and 60/066,029, which have been incorporated by reference in the present application. Preferably, the subject is a vertebrate, more preferably a mammal, and most preferably a human. In preferred aspects, the exendin or exendin agonist is administered parenterally, more preferably by injection. In a most preferred aspect, the injection is a peripheral injection. Preferably, about 10  $\mu$ g-30  $\mu$ g to about 5 mg of the exendin or exendin agonist is administered per day. More preferably, about 10-30  $\mu$ g to about 2mg, or about 10-30  $\mu$ g to about 1mg of the exendin or

exendin agonist is administered per day. Most preferably, about 30  $\mu$ g to about 500  $\mu$ g of the exendin or exendin agonist is administered per day.

5 In various preferred embodiments of the invention, the condition or disorder is obesity, diabetes, preferably Type II diabetes, an eating disorder, or insulin-resistance syndrome.

10 In other preferred aspects of the invention, a method is provided for reducing the appetite of a subject comprising administering to said subject an appetite-lowering amount of an exendin or an exendin agonist.

15 In yet other preferred aspects, a method is provided for lowering plasma lipids comprising administering to said subject a therapeutically effective amount of an exendin or an exendin agonist.

20 The methods of the present invention may also be used to reduce the cardiac risk of a subject comprising administering to said subject a therapeutically effective amount of an exendin or an exendin agonist. In one preferred aspect, the exendin or exendin agonist used in the methods of the present invention is exendin-3. In another preferred aspect, said exendin is exendin-4. Other preferred exendin agonists include exendin-4 (1-30) [SEQ ID NO 6: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln  
25 Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly], exendin-4 (1-30) amide [SEQ ID NO 7: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly-NH<sub>2</sub>], exendin-4 (1-28) amide [SEQ ID NO 40: His Gly Glu Gly Thr

Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg  
Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>], <sup>14</sup>Leu, <sup>25</sup>Phe exendin-4  
amide [SEQ ID NO 9: His Gly Glu Gly Thr Phe Thr Ser Asp Leu  
Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe  
5 Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser-  
NH<sub>2</sub>], <sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 (1-28) amide [SEQ ID NO 41: His  
Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu  
Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>], and  
<sup>14</sup>Leu, <sup>22</sup>Ala, <sup>25</sup>Phe exendin-4 (1-28) amide [SEQ ID NO 8: His Gly  
10 Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu  
Ala Val Arg Leu Ala Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>].

In the methods of the present invention, the exendins  
and exendin agonists may be administered separately or  
together with one or more other compounds and compositions  
15 that exhibit a long term or short-term satiety action,  
including, but not limited to other compounds and  
compositions that comprise an amylin agonist,  
cholecystokinin (CCK), or a leptin (ob protein). Suitable  
amylin agonists include, for example, [<sup>25,28,29</sup>Pro-]-human  
20 amylin (also known as "pramlintide," and previously  
referred to as "AC-137") as described in "Amylin Agonist  
Peptides and Uses Therefor," U.S. Patent No. 5,686,511,  
issued November 11, 1997, and salmon calcitonin. The CCK  
used is preferably CCK octopeptide (CCK-8). Leptin is  
25 discussed in, for example, Pelleymounter, M.A., et al.  
Science 269:540-43 (1995); Halaas, J.L., et al. Science  
269:543-46 (1995); and Campfield, L.A., et al. Eur. J.  
Pharmac. 262:133-41 (1994).

In other embodiments of the invention is provided a pharmaceutical composition for use in the treatment of conditions or disorders which can be alleviated by reducing food intake comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier. Preferably, the pharmaceutical composition comprises a therapeutically effective amount for a human subject.

The pharmaceutical composition may preferably be used for reducing the appetite of a subject, reducing the weight of a subject, lowering the plasma lipid level of a subject, or reducing the cardiac risk of a subject. Those of skill in the art will recognize that the pharmaceutical composition will preferably comprise a therapeutically effective amount of an exendin or exendin agonist to accomplish the desired effect in the subject.

The pharmaceutical compositions may further comprise one or more other compounds and compositions that exhibit a long-term or short-term satiety action, including, but not limited to other compounds and compositions that comprise an amylin agonist, CCK, preferably CCK-8, or leptin. Suitable amylin agonists include, for example, [<sup>25,28,29</sup>Pro]-human amylin and salmon calcitonin.

In one preferred aspect, the pharmaceutical composition comprises exendin-3. In another preferred aspect, the pharmaceutical composition comprises exendin-4.

In other preferred aspects, the pharmaceutical compositions comprises a peptide selected from: exendin-4 (1-30), exendin-4 (1-30) amide, exendin-4 (1-28) amide,



<sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 amide, <sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 (1-28) amide, and <sup>14</sup>Leu, <sup>22</sup>Ala, <sup>25</sup>Phe exendin-4 (1-28) amide.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of exendin-4 and GLP-1.

10

Figure 2 is a graphical depiction of the change of food intake in obese mice after intraperitoneal injection of exendin-4.

Figure 3 is a graphical depiction of the change of food intake in rats after intracerebroventricular injection of exendin-4

15

Figure 4 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of exendin-4 (1-30) ("Compound 1").

20

Figure 5 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of exendin-4 (1-30) amide ("Compound 2").

Figure 6 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of exendin-4 (1-28) amide ("Compound 3").

25

Figure 7 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of <sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 amide ("Compound 4").

Figure 8 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of <sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 (1-28) amide ("Compound 5").

Figure 9 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of <sup>14</sup>Leu, <sup>22</sup>Ala, <sup>25</sup>Phe exendin-4 (1-28) amide ("Compound 6").

Figure 10 depicts the amino acid sequences for certain exendin agonist compounds useful in the present invention [SEQ ID NOS 9-39].

#### DETAILED DESCRIPTION OF THE INVENTION

Exendins and exendin agonists are useful as described herein in view of their pharmacological properties. Activity as exendin agonists can be indicated by activity in the assays described below. Effects of exendins or exendin agonists on reducing food intake can be identified, evaluated, or screened for, using the methods described in the Examples below, or other methods known in the art for determining effects on food intake or appetite.

#### Exendin Agonist Compounds

Exendin agonist compounds are those described in U.S. Provisional Application No. 60/055,404, including compounds of the formula (I) [SEQ ID NO. 3]:

1	5	10
Xaa <sub>1</sub>	Xaa <sub>2</sub>	Xaa <sub>3</sub>
Gly	Thr	Xaa <sub>4</sub>
Xaa <sub>5</sub>	Xaa <sub>6</sub>	Xaa <sub>7</sub>
Xaa <sub>8</sub>	Xaa <sub>9</sub>	Xaa <sub>10</sub>
Ser	Lys	Gln
Xaa <sub>11</sub>	Glu	Glu
Glu	Ala	Val
Arg	Leu	
Xaa <sub>12</sub>	Xaa <sub>13</sub>	Leu
Lys	Asn	Gly
Gly	Xaa <sub>14</sub>	
Ser	Ser	Gly
Ala	Xaa <sub>15</sub>	Xaa <sub>16</sub>
Xaa <sub>17</sub>	Xaa <sub>18</sub>	-Z

wherein Xaa<sub>1</sub> is His, Arg or Tyr; Xaa<sub>2</sub> is Ser, Gly, Ala or Thr; Xaa<sub>3</sub> is Asp or Glu; Xaa<sub>4</sub> is Phe, Tyr or naphthylalanine; Xaa<sub>5</sub> is Thr or Ser; Xaa<sub>6</sub> is Ser or Thr; Xaa<sub>7</sub> is Asp or Glu; Xaa<sub>8</sub> is Leu, Ile, Val, pentylglycine or Met; Xaa<sub>9</sub> is Leu, Ile, pentylglycine, Val or Met; Xaa<sub>10</sub> is Phe, Tyr or naphthylalanine; Xaa<sub>11</sub> is Ile, Val, Leu, pentylglycine, tert-butylglycine or Met; Xaa<sub>12</sub> is Glu or Asp; Xaa<sub>13</sub> is Trp, Phe, Tyr, or naphthylalanine; Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub> and Xaa<sub>17</sub> are independently Pro, homoproline, 3Hyp, 4Hyp, thioproline, N-alkylglycine, N-alkylpentylglycine or N-alkylalanine; Xaa<sub>18</sub> is Ser, Thr or Tyr; and Z is -OH or -NH<sub>2</sub>; with the proviso that the compound is not exendin-3 or exindin-4.

Preferred N-alkyl groups for N-alkylglycine, N-alkylpentylglycine and N-alkylalanine include lower alkyl groups preferably of 1 to about 6 carbon atoms, more preferably of 1 to 4 carbon atoms. Suitable compounds include those listed in Figure 10 having amino acid sequences of SEQ. ID. NOS. 9 to 39.

Preferred exendin agonist compounds include those wherein Xaa<sub>1</sub> is His or Tyr. More preferably Xaa<sub>1</sub> is His.

Preferred are those compounds wherein Xaa<sub>2</sub> is Gly.

Preferred are those compounds wherein Xaa<sub>3</sub> is Leu, pentylglycine or Met.

Preferred compounds include those wherein Xaa<sub>11</sub> is Trp or Phe.

Also preferred are compounds where Xaa<sub>4</sub> is Phe or naphthylalanine; Xaa<sub>11</sub> is Ile or Val and Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>

and Xaa<sub>1</sub>, are independently selected from Pro, homoproline, thioproline or N-alkylalanine. Preferably N-alkylalanine has a N-alkyl group of 1 to about 6 carbon atoms.

According to an especially preferred aspect, Xaa<sub>15</sub>,  
5 Xaa<sub>16</sub> and Xaa<sub>17</sub> are the same amino acid residue.

Preferred are compounds wherein Xaa<sub>18</sub> is Ser or Tyr, more preferably Ser.

Preferably Z is -NH<sub>2</sub>.

According to one aspect, preferred are compounds of  
10 formula (I) wherein Xaa<sub>1</sub> is His or Tyr, more preferably His; Xaa<sub>2</sub> is Gly; Xaa<sub>4</sub> is Phe or naphthylalanine; Xaa<sub>5</sub> is Leu, pentylglycine or Met; Xaa<sub>10</sub> is Phe or naphthylalanine; Xaa<sub>11</sub> is Ile or Val; Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub> and Xaa<sub>17</sub> are  
independently selected from Pro, homoproline, thioproline  
15 or N-alkylalanine; and Xaa<sub>18</sub> is Ser or Tyr, more preferably Ser. More preferably Z is -NH<sub>2</sub>.

According to an especially preferred aspect,  
especially preferred compounds include those of formula  
(I) wherein: Xaa<sub>1</sub> is His or Arg; Xaa<sub>2</sub> is Gly; Xaa<sub>3</sub> is Asp  
20 or Glu; Xaa<sub>4</sub> is Phe or naphthylalanine; Xaa<sub>5</sub> is Thr or Ser;

Xaa<sub>6</sub> is Ser or Thr; Xaa<sub>7</sub> is Asp or Glu; Xaa<sub>8</sub> is Leu or pentylglycine; Xaa<sub>9</sub> is Leu or pentylglycine; Xaa<sub>10</sub> is Phe or naphthylalanine; Xaa<sub>11</sub> is Ile, Val or t-butyltylglycine; Xaa<sub>12</sub> is Glu or Asp; Xaa<sub>13</sub> is Trp or Phe; Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>,  
25 and Xaa<sub>17</sub> are independently Pro, homoproline, thioproline, or N-methylalanine; Xaa<sub>18</sub> is Ser or Tyr; and Z is -OH or -NH<sub>2</sub>; with the proviso that the compound does not have the formula of either SEQ. ID. NOS. 1 or 2. More preferably Z

is -NH<sub>2</sub>. Especially preferred compounds include those having the amino acid sequence of SEQ. ID. NOS. 9, 10, 21, 22, 23, 26, 28, 34, 35 and 39.

5 According to an especially preferred aspect, provided are compounds where Xaa<sub>1</sub> is Leu, Ile, Val or pentylglycine, more preferably Leu or pentylglycine, and Xaa<sub>11</sub> is Phe, Tyr or naphthylalanine, more preferably Phe or naphthylalanine. These compounds will exhibit advantageous  
10 duration of action and be less subject to oxidative degradation, both in vitro and in vivo, as well as during synthesis of the compound.

Exendin agonist compounds also include those described in U.S. Provisional Application No. 60/065,442,  
15 including compounds of the formula (II) [SEQ ID NO. 4]:

Xaa<sub>1</sub> Xaa<sub>2</sub> Xaa<sub>3</sub> Gly Xaa<sub>5</sub> Xaa<sub>6</sub> Xaa<sub>7</sub> Xaa<sub>8</sub> Xaa<sub>9</sub> Xaa<sub>10</sub>  
Xaa<sub>11</sub> Xaa<sub>12</sub> Xaa<sub>13</sub> Xaa<sub>14</sub> Xaa<sub>15</sub> Xaa<sub>16</sub> Xaa<sub>17</sub> Ala Xaa<sub>19</sub> Xaa<sub>20</sub>  
Xaa<sub>21</sub> Xaa<sub>22</sub> Xaa<sub>23</sub> Xaa<sub>24</sub> Xaa<sub>25</sub> Xaa<sub>26</sub> Xaa<sub>27</sub> Xaa<sub>28</sub>-Z<sub>1</sub>; wherein

20

Xaa<sub>1</sub> is His, Arg or Tyr;

Xaa<sub>2</sub> is Ser, Gly, Ala or Thr;

Xaa<sub>3</sub> is Asp or Glu;

Xaa<sub>5</sub> is Ala or Thr;

25

Xaa<sub>6</sub> is Ala, Phe, Tyr or naphthylalanine;

Xaa<sub>7</sub> is Thr or Ser;

Xaa<sub>8</sub> is Ala, Ser or Thr;

Xaa<sub>9</sub> is Asp or Glu;

- Xaa<sub>10</sub> is Ala, Leu, Ile, Val, pentylglycine or Met;  
 Xaa<sub>11</sub> is Ala or Ser;  
 Xaa<sub>12</sub> is Ala or Lys;  
 Xaa<sub>13</sub> is Ala or Gln;
- 5 Xaa<sub>14</sub> is Ala, Leu, Ile, pentylglycine, Val or Met;  
 Xaa<sub>15</sub> is Ala or Glu;  
 Xaa<sub>16</sub> is Ala or Glu;  
 Xaa<sub>17</sub> is Ala or Glu;  
 Xaa<sub>18</sub> is Ala or Val;
- 10 Xaa<sub>20</sub> is Ala or Arg;  
 Xaa<sub>21</sub> is Ala or Leu;  
 Xaa<sub>22</sub> is Ala, Phe, Tyr or naphthylalanine;  
 Xaa<sub>23</sub> is Ile, Val, Leu, pentylglycine, tert-butylglycine  
 or Met;
- 15 Xaa<sub>24</sub> is Ala, Glu or Asp;  
 Xaa<sub>25</sub> is Ala, Trp, Phe, Tyr or naphthylalanine;  
 Xaa<sub>26</sub> is Ala or Leu;  
 Xaa<sub>27</sub> is Ala or Lys;  
 Xaa<sub>28</sub> is Ala or Asn;
- 20 Z<sub>1</sub> is -OH,  
       -NH<sub>2</sub>  
       Gly-Z<sub>2</sub>,  
       Gly Gly-Z<sub>2</sub>,  
       Gly Gly Xaa<sub>31</sub>-Z<sub>2</sub>,
- 25 Gly Gly Xaa<sub>31</sub> Ser-Z<sub>2</sub>,  
       Gly Gly Xaa<sub>31</sub> Ser Ser-Z<sub>2</sub>,  
       Gly Gly Xaa<sub>31</sub> Ser Ser Gly-Z<sub>2</sub>,  
       Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub>-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub>-Z<sub>2</sub> or

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>39</sub>-Z<sub>2</sub>;

Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>39</sub> are independently Pro,

5 homoproline, 3Hyp, 4Hyp, thioproline,

N-alkylglycine, N-alkylpentylglycine or

N-alkylalanine; and

Z<sub>2</sub> is -OH or -NH<sub>2</sub>;

provided that no more than three of Xaa<sub>3</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>, Xaa<sub>8</sub>,

10 Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>, Xaa<sub>17</sub>, Xaa<sub>19</sub>,

Xaa<sub>22</sub>, Xaa<sub>23</sub>, Xaa<sub>24</sub>, Xaa<sub>25</sub>, Xaa<sub>26</sub>, Xaa<sub>27</sub> and Xaa<sub>28</sub> are Ala.

Preferred N-alkyl groups for N-alkylglycine, N-

alkylpentylglycine and N-alkylalanine include lower alkyl

groups preferably of 1 to about 6 carbon atoms, more

15 preferably of 1 to 4 carbon atoms.

Preferred exendin agonist compounds include those  
wherein Xaa<sub>1</sub> is His or Tyr. More preferably Xaa<sub>1</sub> is His.

Preferred are those compounds wherein Xaa<sub>2</sub> is Gly.

Preferred are those compounds wherein Xaa<sub>14</sub> is Leu,  
20 pentylglycine or Met.

Preferred compounds are those wherein Xaa<sub>25</sub> is Trp or  
Phe.

Preferred compounds are those where Xaa<sub>6</sub> is Phe or  
naphthylalanine; Xaa<sub>22</sub> is Phe or naphthylalanine and  
25 Xaa<sub>23</sub> is Ile or Val.

Preferred are compounds wherein Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and  
Xaa<sub>39</sub> are independently selected from Pro, homoproline,  
thioproline and N-alkylalanine.

Preferably  $Z_1$  is  $-NH_2$ .

Preferable  $Z_2$  is  $-NH_2$ .

According to one aspect, preferred are compounds of formula (I) wherein  $Xaa_1$  is His or Tyr, more preferably His;  $Xaa_2$  is Gly;  $Xaa_3$  is Phe or naphthylalanine;  $Xaa_{14}$  is Leu, pentylglycine or Met;  $Xaa_{22}$  is Phe or naphthylalanine;  $Xaa_{23}$  is Ile or Val;  $Xaa_{31}$ ,  $Xaa_{36}$ ,  $Xaa_{37}$  and  $Xaa_{38}$  are independently selected from Pro, homoproline, thioproline or N-alkylalanine. More preferably  $Z_1$  is  $-NH_2$ .

According to an especially preferred aspect, especially preferred compounds include those of formula (I) wherein:  $Xaa_1$  is His or Arg;  $Xaa_2$  is Gly or Ala;  $Xaa_3$  is Asp or Glu;  $Xaa_4$  is Ala or Thr;  $Xaa_5$  is Ala, Phe or naphthylalanine;  $Xaa_6$  is Thr or Ser;  $Xaa_8$  is Ala, Ser or Thr;  $Xaa_9$  is Asp or Glu;  $Xaa_{10}$  is Ala, Leu or pentylglycine;  $Xaa_{11}$  is Ala or Ser;  $Xaa_{12}$  is Ala or Lys;  $Xaa_{13}$  is Ala or Gln;  $Xaa_{14}$  is Ala, Leu or pentylglycine;  $Xaa_{15}$  is Ala or Glu;  $Xaa_{16}$  is Ala or Glu;  $Xaa_{17}$  is Ala or Glu;  $Xaa_{19}$  is Ala or Val;  $Xaa_{20}$  is Ala or Arg;  $Xaa_{21}$  is Ala or Leu;  $Xaa_{22}$  is Phe or naphthylalanine;  $Xaa_{23}$  is Ile, Val or tert-butylglycine;  $Xaa_{24}$  is Ala, Glu or Asp;  $Xaa_{25}$  is Ala, Trp or Phe;  $Xaa_{26}$  is Ala or Leu;  $Xaa_{27}$  is Ala or Lys;  $Xaa_{28}$  is Ala or Asn;  $Z_1$  is  $-OH$ ,  $-NH_2$ , Gly- $Z_2$ , Gly Gly- $Z_2$ , Gly Gly  $Xaa_{31}$ - $Z_2$ , Gly Gly  $Xaa_{31}$  Ser- $Z_2$ , Gly Gly  $Xaa_{31}$  Ser Ser- $Z_2$ , Gly Gly  $Xaa_{31}$  Ser Ser Gly- $Z_2$ , Gly Gly  $Xaa_{31}$  Ser Ser Gly Ala- $Z_2$ , Gly Gly  $Xaa_{31}$  Ser Ser Gly Ala  $Xaa_{36}$ - $Z_2$ , Gly Gly  $Xaa_{31}$  Ser Ser Gly Ala  $Xaa_{36}$   $Xaa_{37}$ - $Z_2$ , Gly Gly  $Xaa_{31}$  Ser Ser Gly Ala  $Xaa_{36}$   $Xaa_{37}$   $Xaa_{38}$ - $Z_2$ ;  $Xaa_{31}$ ,  $Xaa_{36}$ ,  $Xaa_{37}$  and  $Xaa_{38}$  being independently Pro homoproline, thioproline or N-



methylaniline; and  $Z_2$  being -OH or -NH<sub>2</sub>; provided that no more than three of Xaa<sub>3</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>, Xaa<sub>8</sub>, Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>, Xaa<sub>17</sub>, Xaa<sub>19</sub>, Xaa<sub>20</sub>, Xaa<sub>21</sub>, Xaa<sub>24</sub>, Xaa<sub>25</sub>, Xaa<sub>26</sub>, Xaa<sub>27</sub> and Xaa<sub>28</sub> are Ala. Especially  
 5 preferred compounds include those having the amino acid sequence of SEQ. ID. NOS. 40-61.

According to an especially preferred aspect, provided are compounds where Xaa<sub>4</sub> is Leu, Ile, Val or pentylglycine, more preferably Leu or pentylglycine, and  
 10 Xaa<sub>25</sub> is Phe, Tyr or naphthylalanine, more preferably Phe or naphthylalanine. These compounds will be less susceptible to oxidative degradation, both in vitro and in vivo, as well as during synthesis of the compound.

Exendin agonist compounds also include those  
 15 described in U.S. Provisional Application No. 60/066,029, including compounds of the formula (III) [SEQ ID NO. 5]:

Xaa<sub>1</sub> Xaa<sub>2</sub> Xaa<sub>3</sub> Xaa<sub>4</sub> Xaa<sub>5</sub> Xaa<sub>6</sub> Xaa<sub>7</sub> Xaa<sub>8</sub> Xaa<sub>9</sub> Xaa<sub>10</sub>  
 Xaa<sub>11</sub> Xaa<sub>12</sub> Xaa<sub>13</sub> Xaa<sub>14</sub> Xaa<sub>15</sub> Xaa<sub>16</sub> Xaa<sub>17</sub> Ala Xaa<sub>19</sub> Xaa<sub>20</sub>  
 20 Xaa<sub>21</sub> Xaa<sub>22</sub> Xaa<sub>23</sub> Xaa<sub>24</sub> Xaa<sub>25</sub> Xaa<sub>26</sub> Xaa<sub>27</sub> Xaa<sub>28</sub>-Z<sub>1</sub>; wherein

Xaa<sub>1</sub> is His, Arg, Tyr, Ala, Norval, Val or Norleu;  
 Xaa<sub>2</sub> is Ser, Gly, Ala or Thr;  
 25 Xaa<sub>3</sub> is Ala, Asp or Glu;  
 Xaa<sub>4</sub> is Ala, Norval, Val, Norleu or Gly;  
 Xaa<sub>5</sub> is Ala or Thr;  
 Xaa<sub>6</sub> is Phe, Tyr or naphthylalanine;

- Xaa<sub>1</sub> is Thr or Ser;
- Xaa<sub>2</sub> is Ala, Ser or Thr;
- Xaa<sub>3</sub> is Ala, Norval, Val, Norleu, Asp or Glu;
- Xaa<sub>10</sub> is Ala, Leu, Ile, Val, pentylglycine or Met;
- 5 Xaa<sub>11</sub> is Ala or Ser;
- Xaa<sub>12</sub> is Ala or Lys;
- Xaa<sub>13</sub> is Ala or Gln;
- Xaa<sub>14</sub> is Ala, Leu, Ile, pentylglycine, Val or Met;
- Xaa<sub>15</sub> is Ala or Glu;
- 10 Xaa<sub>16</sub> is Ala or Glu;
- Xaa<sub>17</sub> is Ala or Glu;
- Xaa<sub>19</sub> is Ala or Val;
- Xaa<sub>20</sub> is Ala or Arg;
- Xaa<sub>21</sub> is Ala or Leu;
- 15 Xaa<sub>22</sub> is Phe, Tyr or naphthylalanine;
- Xaa<sub>23</sub> is Ile, Val, Leu, pentylglycine, tert-butylglycine or Met;
- Xaa<sub>24</sub> is Ala, Glu or Asp;
- Xaa<sub>25</sub> is Ala, Trp, Phe, Tyr or naphthylalanine;
- 20 Xaa<sub>26</sub> is Ala or Leu;
- Xaa<sub>27</sub> is Ala or Lys;
- Xaa<sub>28</sub> is Ala or Asn;
- Z<sub>1</sub> is -OH,  
           -NH<sub>2</sub>,
- 25 Gly-Z<sub>2</sub>,
- Gly Gly-Z<sub>2</sub>,
- Gly Gly Xaa<sub>31</sub>-Z<sub>2</sub>,
- Gly Gly Xaa<sub>31</sub> Ser-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser-Z<sub>2</sub>,  
 Gly Gly Xaa<sub>31</sub> Ser Ser Gly-Z<sub>2</sub>,  
 Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala-Z<sub>2</sub>,  
 Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub>-Z<sub>2</sub>,  
 5 Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub>-Z<sub>2</sub>,  
 Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub>-Z<sub>2</sub> or Gly  
 Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub> Xaa<sub>39</sub>-Z<sub>2</sub>; wherein  
 Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub>, and Xaa<sub>38</sub> are independently  
 Pro, homoproline, 3Hyp, 4Hyp, thioproline,  
 10 N-alkylglycine, N-alkylpentylglycine or  
 N-alkylalanine; and  
 Z<sub>2</sub> is -OH or -NH<sub>2</sub>;

provided that no more than three of Xaa<sub>3</sub>, Xaa<sub>4</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>,  
 Xaa<sub>8</sub>, Xaa<sub>9</sub>, Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>,  
 15 Xaa<sub>17</sub>, Xaa<sub>19</sub>, Xaa<sub>20</sub>, Xaa<sub>21</sub>, Xaa<sub>24</sub>, Xaa<sub>25</sub>, Xaa<sub>26</sub>, Xaa<sub>27</sub> and Xaa<sub>28</sub>  
 are Ala; and provided also that, if Xaa<sub>1</sub> is His, Arg or  
 Tyr, then at least one of Xaa<sub>3</sub>, Xaa<sub>4</sub> and Xaa<sub>9</sub> is Ala.

### Definitions

20 In accordance with the present invention and as used  
 herein, the following terms are defined to have the  
 following meanings, unless explicitly stated otherwise.


The term "amino acid" refers to natural amino acids,  
 unnatural amino acids, and amino acid analogs, all in  
 25 their D and L stereoisomers if their structure allow such  
 stereoisomeric forms. Natural amino acids include alanine  
 (Ala), arginine (Arg), asparagine (Asn), aspartic acid  
 (Asp), cysteine (Cys), glutamine (Gln), glutamic acid  
 (Glu), glycine (Gly), histidine (His), isoleucine (Ile),

leucine (Leu), Lysine (Lys), methionine (Met),  
phenylalanine (Phe), proline (Pro), serine (Ser),  
threonine (Thr), typtophan (Trp), tyrosine (Tyr) and  
valine (Val). Unnatural amino acids include, but are not  
5 limited to azetidinecarboxylic acid, 2-aminoadipic acid,  
3-aminoadipic acid, beta-alanine, aminopropionic acid, 2-  
aminobutyric acid, 4-aminobutyric acid, 6-aminocaproic  
acid, 2-aminoheptanoic acid, 2-aminoisobutyric acid, 3-  
aminoisbutyric acid, 2-aminopimelic acid, tertiary-  
10 butylglycine, 2,4-diaminoisobutyric acid, desmosine, 2,2'-  
diaminopimelic acid, 2,3-diaminopropionic acid, N-  
ethylglycine, N-ethylasparagine, homoproline,  
hydroxylysine, allo-hydroxylysine, 3-hydroxyproline, 4-  
hydroxyproline, isodesmosine, allo-isoleucine, N-  
15 methylalanine, N-methylglycine, N-methylisoleucine, N-  
methylpentylglycine, N-methylvaline, naphthalanine,  
norvaline, norleucine, ornithine, pentylglycine, pipercolic  
acid and thioproline. Amino acid analogs include the  
natural and unnatural amino acids which are chemically  
20 blocked, reversibly or irreversibly, or modified on their  
N-terminal amino group or their side-chain groups, as for  
example, methionine sulfoxide, methionine sulfone, S-  
(carboxymethyl)-cysteine, S-(carboxymethyl)-cysteine  
sulfoxide and S-(carboxymethyl)-cysteine sulfone.

25 The term "amino acid analog" refers to an amino acid  
wherein either the C-terminal carboxy group, the N-  
terminal amino group or side-chain functional group has  
been chemically codified to another functional group. For  
example, aspartic acid-(beta-methyl ester) is an amino

acid analog of aspartic acid; N-ethylglycine is an amino acid analog of glycine; or alanine carboxamide is an amino acid analog of alanine.

The term "amino acid residue" refers to radicals  
 5 having the structure: (1)  $-C(O)-R-NH-$ , wherein R typically is  $-CH(R')$ -, wherein R' is an amino acid side chain, typically H or a carbon containing substituent; or (2),



10

wherein p is 1, 2 or 3 representing the azetidinecarboxylic acid, proline or pipercolic acid residues, respectively.

The term "lower" referred to herein in connection  
 15 with organic radicals such as alkyl groups defines such groups with up to and including about 6, preferably up to and including 4 and advantageously one or two carbon atoms. Such groups may be straight chain or branched chain.

20 "Pharmaceutically acceptable salt" includes salts of the compounds described herein derived from the combination of such compounds and an organic or inorganic acid. In practice the use of the salt form amounts to use of the base form. The compounds are useful in both free  
 25 base and salt form.

In addition, the following abbreviations stand for the following:

"ACN" or "CH<sub>3</sub>CN" refers to acetonitrile.

"Boc", "tBoc" or "Tboc" refers to t-butoxy carbonyl.

5 "DCC" refers to N,N'-dicyclohexylcarbodiimide.

"Fmoc" refers to fluorenylmethoxycarbonyl.

"HBTU" refers to 2-(1H-benzotriazol-1-yl)-

1,1,3,3,-tetramethyluronium hexafluorophosphate.

"HOBt" refers to 1-hydroxybenzotriazole monohydrate.

10 "homoP" or hPro" refers to homoproline.

"MeAla" or "Nme" refers to N-methylalanine.

"naph" refers to naphthylalanine.

"pG" or pGly" refers to pentylglycine.

"tBuG" refers to tertiary-butylglycine.

15 "ThioP" or tPro" refers to thioproline.

3Hyp" refers to 3-hydroxyproline

4Hyp" refers to 4-hydroxyproline.

NAG" refers to N-alkylglycine

NAPG" refers to N-alkylpentylglycine

20 "Norval" refers to norvaline

"Norleu" refers to norleucine

#### Preparation of Compounds

25 The exendins and exendin agonists described herein may be prepared using standard solid-phase peptide synthesis techniques and preferably an automated or semiautomated peptide synthesizer. Typically, using such

techniques, an  $\alpha$ -N-carbamoyl protected amino acid and an amino acid attached to the growing peptide chain on a resin are coupled at room temperature in an inert solvent such as dimethylformamide, N-methylpyrrolidinone or methylene chloride in the presence of coupling agents such as dicyclohexylcarbodiimide and 1-hydroxybenzotriazole in the presence of a base such as diisopropylethylamine. The  $\alpha$ -N-carbamoyl protecting group is removed from the resulting peptide-resin using a reagent such as trifluoroacetic acid or piperidine, and the coupling reaction repeated with the next desired N-protected amino acid to be added to the peptide chain. Suitable N-protecting groups are well known in the art, with t-butyloxycarbonyl (tBoc) and fluorenylmethoxycarbonyl (Fmoc) being preferred herein.

The solvents, amino acid derivatives and 4-methylbenzhydryl-amine resin used in the peptide synthesizer may be purchased from Applied Biosystems Inc. (Foster City, CA). The following side-chain protected amino acids may be purchased from Applied Biosystems, Inc.: Boc-Arg(Mts), Fmoc-Arg(Pmc), Boc-Thr(Bzl), Fmoc-Thr(t-Bu), Boc-Ser(Bzl), Fmoc-Ser(t-Bu), Boc-Tyr(BrZ), Fmoc-Tyr(t-Bu), Boc-Lys(Cl-Z), Fmoc-Lys(Boc), Boc-Glu(Bzl), Fmoc-Glu(t-Bu), Fmoc-His(Trt), Fmoc-Asn(Trt), and Fmoc-Gln(Trt). Boc-His(BOM) may be purchased from Applied Biosystems, Inc. or Bachem Inc. (Torrance, CA). Anisole, dimethylsulfide, phenol, ethanedithiol, and thioanisole may be obtained from

Aldrich Chemical Company (Milwaukee, WI). Air Products and Chemicals (Allentown, PA) supplies HF. Ethyl ether, acetic acid and methanol may be purchased from Fisher Scientific (Pittsburgh, PA).

5           Solid phase peptide synthesis may be carried out with an automatic peptide synthesizer (Model 430A, Applied Biosystems Inc., Foster City, CA) using the NMP/HOBt (Option 1) system and tBoc or Fmoc chemistry (see, Applied Biosystems User's Manual for the ABI 430A Peptide  
10           Synthesizer, Version 1.3B July 1, 1988, section 6, pp. 49-70, Applied Biosystems, Inc., Foster City, CA) with capping. Boc-peptide-resins may be cleaved with HF (-5° C to 0° C, 1 hour). The peptide may be extracted from the resin with alternating water and acetic acid, and the  
15           filtrates lyophilized. The Fmoc-peptide resins may be cleaved according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems, Inc., 1990, pp. 6-12). Peptides may be also be assembled using an  
20           Advanced Chem Tech Synthesizer (Model MPS 350, Louisville, Kentucky).

          Peptides may be purified by RP-HPLC (preparative and analytical) using a Waters Delta Prep 3000 system. A C4, C8 or C18 preparative column (10 $\mu$  , 2.2 x 25 cm; Vydac, Hesperia, CA) may be used to isolate peptides, and purity  
25           may be determined using a C4, C8 or C18 analytical column (5 $\mu$  , 0.46 x 25 cm; Vydac). Solvents (A=0.1% TFA/water and B=0.1% TFA/CH<sub>3</sub>CN) may be delivered to the analytical column at a flowrate of 1.0 ml/min and to the preparative column at 15 ml/min. Amino acid analyses may be performed



on the Waters Pico Tag system and processed using the Maxima program. Peptides may be hydrolyzed by vapor-phase acid hydrolysis (115° C, 20-24 h). Hydrolysates may be derivatized and analyzed by standard methods (Cohen, et al., The Pico Tag Method: A Manual of Advanced Techniques for Amino Acid Analysis, pp. 11-52, Millipore Corporation, Milford, MA (1989)). Fast atom bombardment analysis may be carried out by M-Scan, Incorporated (West Chester, PA).

Mass calibration may be performed using cesium iodide or cesium iodide/glycerol. Plasma desorption ionization analysis using time of flight detection may be carried out on an Applied Biosystems Bio-Ion 20 mass spectrometer. Electrospray mass spectroscopy may be carried out on a VG-Trio machine.

Peptide compounds useful in the invention may also be prepared using recombinant DNA techniques, using methods now known in the art. See, e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2d Ed., Cold Spring Harbor (1989). Non-peptide compounds useful in the present invention may be prepared by art-known methods. For example, phosphate-containing amino acids and peptides containing such amino acids, may be prepared using methods known in the art. See, e.g., Bartlett and Landen, Biorq. Chem. 14:356-377 (1986).

The compounds described above are useful in view of their pharmacological properties. In particular, the compounds of the invention possess activity as agents to reduce food intake. They can be used to treat conditions or diseases which can be alleviated by reducing food

intake.

Compositions useful in the invention may conveniently be provided in the form of formulations suitable for parenteral (including intravenous, intramuscular and subcutaneous) or nasal or oral administration. In some cases, it will be convenient to provide an exendin or exendin agonist and another food-intake-reducing, plasma glucose-lowering or plasma lipid-lowering agent, such as amylin, an amylin agonist, a CCK, or a leptin, in a single composition or solution for administration together. In other cases, it may be more advantageous to administer the additional agent separately from said exendin or exendin agonist. A suitable administration format may best be determined by a medical practitioner for each patient individually. Suitable pharmaceutically acceptable carriers and their formulation are described in standard formulation treatises, e.g., Remington's Pharmaceutical Sciences by E.W. Martin. See also Wang, Y.J. and Hanson, M.A. "Parenteral Formulations of Proteins and Peptides: Stability and Stabilizers," Journal of Parenteral Science and Technology, Technical Report No. 10, Supp. 42:2S (1988).

Compounds useful in the invention can be provided as parenteral compositions for injection or infusion. They can, for example, be suspended in an inert oil, suitably a vegetable oil such as sesame, peanut, olive oil, or other acceptable carrier. Preferably, they are suspended in an aqueous carrier, for example, in an isotonic buffer solution at a pH of about 3.0 to 8.0, preferably at a pH of

about 3.5 to 5.0. These compositions may be sterilized by conventional sterilization techniques, or may be sterile filtered. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH buffering agents. Useful buffers include for example, sodium acetate/acetic acid buffers. A form of repository or "depot" slow release preparation may be used so that therapeutically effective amounts of the preparation are delivered into the bloodstream over many hours or days following transdermal injection or delivery.

The desired isotonicity may be accomplished using sodium chloride or other pharmaceutically acceptable agents such as dextrose, boric acid, sodium tartrate, propylene glycol, polyols (such as mannitol and sorbitol), or other inorganic or organic solutes. Sodium chloride is preferred particularly for buffers containing sodium ions.

The claimed compositions can also be formulated as pharmaceutically acceptable salts (e.g., acid addition salts) and/or complexes thereof. Pharmaceutically acceptable salts are non-toxic salts at the concentration at which they are administered. The preparation of such salts can facilitate the pharmacological use by altering the physical-chemical characteristics of the composition without preventing the composition from exerting its physiological effect. Examples of useful alterations in physical properties include lowering the melting point to facilitate transmucosal administration and increasing the solubility to facilitate the administration of higher

concentrations of the drug.

Pharmaceutically acceptable salts include acid addition salts such as those containing sulfate, hydrochloride, phosphate, sulfamate, acetate, citrate, lactate, tartrate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate, cyclohexylsulfamate and quinate. Pharmaceutically acceptable salts can be obtained from acids such as hydrochloric acid, sulfuric acid, phosphoric acid, sulfamic acid, acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, cyclohexylsulfamic acid, and quinic acid. Such salts may be prepared by, for example, reacting the free acid or base forms of the product with one or more equivalents of the appropriate base or acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is then removed in vacuo or by freeze-drying or by exchanging the ions of an existing salt for another ion on a suitable ion exchange resin.

Carriers or excipients can also be used to facilitate administration of the compound. Examples of carriers and excipients include calcium carbonate, calcium phosphate, various sugars such as lactose, glucose, or sucrose, or types of starch, cellulose derivatives, gelatin, vegetable oils, polyethylene glycols and physiologically compatible solvents. The compositions or pharmaceutical composition can be administered by different routes including intravenously, intraperitoneal, subcutaneous, and intramuscular, orally, topically, transmucosally, or by

pulmonary inhalation.

If desired, solutions of the above compositions may be thickened with a thickening agent such as methyl cellulose.

They may be prepared in emulsified form, either water in oil or oil in water. Any of a wide variety of pharmaceutically acceptable emulsifying agents may be employed including, for example, acacia powder, a non-ionic surfactant (such as a Tween), or an ionic surfactant (such as alkali polyether alcohol sulfates or sulfonates, e.g., a Triton).

Compositions useful in the invention are prepared by mixing the ingredients following generally accepted procedures. For example, the selected components may be simply mixed in a blender or other standard device to produce a concentrated mixture which may then be adjusted to the final concentration and viscosity by the addition of water or thickening agent and possibly a buffer to control pH or an additional solute to control tonicity.

For use by the physician, the compositions will be provided in dosage unit form containing an amount of an exendin or exendin agonist, for example, exendin-3, and/or exendin-4, with or without another food intake-reducing, plasma glucose-lowering or plasma lipid-lowering agent. Therapeutically effective amounts of an exendin or exendin agonist for use in reducing food intake are those that suppress appetite at a desired level. As will be recognized by those in the field, an effective amount of therapeutic agent will vary with many factors including the age and weight of the patient, the patient's physical

condition, the blood sugar level and other factors.

The effective daily appetite-suppressing dose of the compounds will typically be in the range of about 10 to 30  $\mu\text{g}$  to about 5 mg/day, preferably about 10 to 30  $\mu\text{g}$  to about 2 mg/day and more preferably about 10 to 100  $\mu\text{g}$  to about 1 mg/day, most preferably about 30  $\mu\text{g}$  to about 500  $\mu\text{g}$ /day, for a 70 kg patient, administered in a single or divided doses. The exact dose to be administered is determined by the attending clinician and is dependent upon where the particular compound lies within the above quoted range, as well as upon the age, weight and condition of the individual. Administration should begin whenever the suppression of food intake, or weight lowering is desired, for example, at the first sign of symptoms or shortly after diagnosis of obesity, diabetes mellitus, or insulin-resistance syndrome. Administration may be by injection, preferably subcutaneous or intramuscular. Orally active compounds may be taken orally, however dosages should be increased 5-10 fold.

The optimal formulation and mode of administration of compounds of the present application to a patient depend on factors known in the art such as the particular disease or disorder, the desired effect, and the type of patient. While the compounds will typically be used to treat human subjects they may also be used to treat similar or identical diseases in other vertebrates such as other primates, farm animals such as swine, cattle and poultry, and sports animals and pets such as horses, dogs and cats.

To assist in understanding the present invention, the following Examples are included. The experiments relating to this invention should not, of course, be construed as specifically limiting the invention and such variations of the invention, now known or later developed, which would be within the purview of one skilled in the art are considered to fall within the scope of the invention as described herein and hereinafter claimed.

EXAMPLE 1: Exendin Injections Reduced the Food Intake of  
Normal Mice

All mice (NIH:Swiss mice) were housed in a stable environment of 22 ( $\pm$  2) $^{\circ}$  C, 60 ( $\pm$ 10) % humidity and a 12:12 light:dark cycle; with lights on at 0600. Mice were housed in groups of four in standard cages with *ad libitum* access to food (Teklad: LM 485; Madison, WI) and water except as noted, for at least two weeks before the experiments.

All experiments were conducted between the hours of 0700 and 0900. The mice were food deprived (food removed at 1600 hr from all animals on day prior to experiment) and individually housed. All mice received an intraperitoneal injection (5  $\mu$ l/kg) of either saline or exendin-4 at doses of 0.1, 1.0, 10 and 100  $\mu$ g/kg and were immediately presented with a pre-weighed food pellet (Teklad LM 485). The food pellet was weighed at 30-minute, 1-hr, 2-hr and 6-hr intervals to determine the amount of food eaten.

Figure 1 depicts cumulative food intake over periods of 0.5, 1, 2 and 6hr in overnight-fasted normal NIH:Swiss

mice following ip injection of saline, 2 doses of GLP-1, or 4 doses of exendin-4. At doses up to 100µg/kg, GLP-1 had no effect on food intake measured over any period, a result consistent with that previously reported (Bhavsar, S.P., et al., Soc. Neurosci. Abstr. 21:460 (188.8) (1995); and Turton, M.D., Nature, 379:69-72, (1996)).

In contrast, exendin-4 injections potently and dose-dependently inhibited food intake. The ED<sub>50</sub> for inhibition of food intake over 30 min was 1µg/kg, which is a level about as potent as amylin (ED<sub>50</sub> 3.6µg/kg) or the prototypical peripheral satiety agent, CCK (ED<sub>50</sub> 0.97µg/kg) as measured in this preparation. However, in contrast to the effects of amylin or CCK, which abate after 1-2 hours, the inhibition of food intake with exendin-4 was still present after at least 6 hours after injection.

#### EXAMPLE 2: Exendin Reduced the Food Intake of Obese Mice

All mice (female ob/ob mice) were housed in a stable environment of 22 (±2)° C; 60 (±10) % humidity and a 12:12 light:dark cycle; with lights on at 0600. Mice were housed in groups of four in standard cages with *ad libitum* access to food (Teklad: LM 485) and water except as noted, for at least two weeks before the experiments.

All experiments were conducted between the hours of 0700 and 0900. The mice were food deprived (food removed at 1600 hr from all animals on day prior to experiment) and individually housed. All mice received an intraperitoneal injection (5 µl/kg) of either saline or exendin-4 at doses



of 0.1, 1.0 and 10  $\mu\text{g/kg}$  (female ob/ob mice) and were immediately presented with a pre-weighed food pellet (Teklad LM 485). The food pellet was weighed at 30-minute, 1 -hr, 2-hr and 6-hr intervals to determine the amount of food eaten.

Figure 2 depicts the effect of exendin-4 in the ob/ob mouse model of obesity. The obese mice had a similar food intake-related response to exendin as the normal mice. Moreover, the obese mice were not hypersensitive to exendin, as has been observed with amylin and leptin (Young, A.A., et al., Program and Abstracts, 10th International Congress of Endocrinology, June 12-15, 1996 San Francisco, pg 419 (P2-58)).

EXAMPLE 3: Intracerebroventricular Injections of Exendin  
Inhibited Food Intake in Rats

All rats (Harlan Sprague-Dawley) were housed in a stable environment of 22' ( $\pm 2$ )° C, 60 ( $\pm 10$ )% humidity and a 12:12 light:dark cycle; with lights on at 0600. Rats were obtained from Zivic Miller with an intracerebroventricular cannula (ICV cannula) implanted (coordinates determined by actual weight of animals and referenced to Paxinos, G. and Watson, C. "The Rat Brain in stereotaxic coordinates," second edition. Academic Press) and were individually housed in standard cages with *ad libitum* access to food (Teklad: LM 485) and water for at least one week before the experiments.

All injections were given between the hours of 1700

and 1800. The rats were habituated to the ICV injection procedure at least once before the ICV administration of compound. All rats received an ICV injection (2  $\mu$ l/30 seconds) of either saline or exendin-4 at doses of 0.01, 0.03, 0.1, 0.3, and 1.0  $\mu$ g. All animals were then presented with pre-weighed food (Teklad LM 485) at 1800, when the lights were turned off. The amount of food left was weighed at 2-hr, 12-hr and 24-hr intervals to determine the amount of food eaten by each animal.

Figure 3 depicts a dose-dependent inhibition of food intake in rats that received doses greater than 0.1 $\mu$ g/rat. The ED<sub>50</sub> was  $\approx$  0.1 $\mu$ g, exendin-4 is thus  $\approx$ 100-fold more potent than intracerebroventricular injections of GLP-1 as reported by Turton, M.D., et al. (Nature 379:69-72 (1996)).

EXAMPLE 4: Exendin Agonists Reduced the Food Intake in Mice

All mice (NIH:Swiss mice) were housed in a stable environment of 22 ( $\pm$ 2)° C, 60 ( $\pm$ 10) % humidity and a 12:12 light:dark cycle; with lights on at 0600. Mice were housed in groups of four in standard cages with *ad libitum* access to food (Teklad: LM 485; Madison, WI) and water except as noted, for at least two weeks before the experiments.

All experiments were conducted between the hours of 0700 and 0900. The mice were food deprived (food removed at 1600 hr from all animals on day prior to experiment) and individually housed. All mice received an

intraperitoneal injection (5  $\mu$ l/kg) of either saline or test compound at doses of 1, 10, and 100  $\mu$ g/kg and immediately presented with a food pellet (Teklad LM 485).

5 The food pellet was weighed at 30-minute, 1-hr, 2-hr and 6-hr intervals to determine the amount of food eaten.

Figure 4 depicts the cumulative food intake over periods of 0.5, 1, 2 and 6 hr in overnight-fasted normal NIH:Swiss mice following ip injection of saline or exendin-4 (1-30) ("Compound 1") in doses of 1, 10 and 100  $\mu$ g/kg.

Figure 5 depicts the cumulative food intake over periods of 0.5, 1, 2 and 6 hr in overnight-fasted normal NIH:Swiss mice following ip injection of saline or exendin-4 (1-30) amide ("Compound 2") in doses of 1, 10 and 100  $\mu$ g/kg.

Figure 6 depicts the cumulative food intake over periods of 0.5, 1, 2 and 6 hr in overnight-fasted normal NIH:Swiss mice following ip injection of saline or exendin-4 (1-28) amide ("Compound 3") in doses of 1, 10 and 100  $\mu$ g/kg.

Figure 7 depicts the cumulative food intake over periods of 0.5, 1, 2 and 6 hr in overnight-fasted normal NIH:Swiss mice following ip injection of saline or <sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 amide ("Compound 4") in doses of 1, 10 and 100  $\mu$ g/kg.

Figure 8 depicts the cumulative food intake over periods of 0.5, 1, 2 and 6 hr in overnight-fasted normal NIH:Swiss mice following ip injection of saline or <sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 (1-28) amide ("Compound 5") in doses

of 1, 10 and 100  $\mu\text{g/kg}$ .

Figure 9 depicts the cumulative food intake over periods of 0.5, 1, 2 and 6 hr in overnight-fasted normal NIH:Swiss mice following ip injection of saline or <sup>14</sup>Leu, <sup>22</sup>Ala, <sup>25</sup>Phe exendin-4 (1-28) amide ("Compound 6") in doses of 1, 10 and 100  $\mu\text{g/kg}$ .

#### EXAMPLE 5

##### Preparation of amidated peptide having SEQ. ID. NO. 9

10

15

20

25

The above-identified peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.). In general, single-coupling cycles were used throughout the synthesis and Fast Moc (HBTU activation) chemistry was employed. However, at some positions coupling was less efficient than expected and double couplings were required. In particular, residues Asp<sub>9</sub>, Thr, and Phe<sub>6</sub> all required double coupling. Deprotection (Fmoc group removal) of the growing peptide chain using piperidine was not always efficient. Double deprotection was required at positions Arg<sub>20</sub>, Val<sub>19</sub>, and Leu<sub>14</sub>. Final deprotection of the completed peptide resin was achieved using a mixture of triethylsilane (0.2 mL), ethanedithiol (0.2 mL), anisole (0.2 mL), water (0.2 mL) and trifluoroacetic acid (15 mL) according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems, Inc.) The peptide was precipitated in ether/water (50 mL) and centrifuged. The

precipitate was reconstituted in glacial acetic acid and lyophilized. The lyophilized peptide was dissolved in water). Crude purity was about 55%.

Used in purification steps and analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).

The solution containing peptide was applied to a preparative C-18 column and purified (10% to 40% Solvent B in Solvent A over 40 minutes). Purity of fractions was determined isocratically using a C-18 analytical column. Pure fractions were pooled furnishing the above-identified peptide. Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.5 minutes. Electrospray Mass Spectrometry (M): calculated 4131.7; found 4129.3.

#### EXAMPLE 6

##### Preparation of Peptide having SEQ. ID. NO. 10

The above-identified peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 25% to 75% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 21.5

minutes. Electrospray Mass Spectrometry (M): calculated 4168.6; found 4171.2.

#### EXAMPLE 7

##### Preparation of Peptide having SEQ. ID. NO. 11

The above-identified peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 17.9 minutes. Electrospray Mass Spectrometry (M): calculated 4147.6; found 4150.2.

#### EXAMPLE 8

##### Preparation of Peptide having SEQ. ID. NO. 12

The above-identified peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).

Analytical RP-HPLC (gradient 35% to 65% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 19.7 minutes. Electrospray Mass Spectrometry (M): calculated 4212.6; found 4213.2.

#### EXAMPLE 9

##### Preparation of Peptide having SEQ. ID. NO. 13

The above-identified peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 50% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 16.3 minutes. Electrospray Mass Spectrometry (M): calculated 4262.7; found 4262.4.

#### EXAMPLE 10

##### Preparation of Peptide having SEQ. ID. NO. 14

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.),

cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4172.6

#### EXAMPLE 11

##### Preparation of Peptide having SEQ. ID. NO. 15

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4224.7.



EXAMPLE 12Preparation of Peptide having SEQ. ID. NO. 16

5           The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
10   similar way to Example 5. Used in analysis are Solvent A  
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
15   product peptide. Electrospray Mass Spectrometry (M):  
calculated 4172.6

EXAMPLE 13Preparation of Peptide having SEQ. ID. NO. 17

20           The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
25   cleaved from the resin, deprotected and purified in a  
similar way to Example 5. Used in analysis are Solvent A  
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is

then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4186.6

5

EXAMPLE 14Preparation of Peptide having SEQ. ID. NO. 18

10

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).

15

Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4200.7

20

EXAMPLE 15Preparation of Peptide having SEQ. ID. NO. 19

25

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis are Solvent A

(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 4200.7

#### EXAMPLE 16

##### Preparation of Peptide having SEQ. ID. NO. 20

The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 5. Used in analysis are Solvent A  
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 4202.7.

#### EXAMPLE 17

##### Preparation of Peptide having SEQ. ID. NO. 21

The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using

Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 5. Used in analysis are Solvent A  
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).

5 Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 4145.6.

10

#### EXAMPLE 18

##### Preparation of Peptide having SEQ. ID. NO. 22

The above-identified peptide is assembled on 4-(2'-  
15 4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 5. Used in analysis are Solvent A  
20 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
25 calculated 4184.6.

EXAMPLE 19Preparation of Peptide having SEQ. ID. NO. 23

5           The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
10       similar way to Example 5. Used in analysis are Solvent A  
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
15       product peptide. Electrospray Mass Spectrometry (M):  
calculated 4145.6.

EXAMPLE 20

20       Preparation of Peptide having SEQ. ID. NO. 24

          The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
25       Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 5. Used in analysis are Solvent A  
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in

Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4224.7.

5

#### EXAMPLE 21

##### Preparation of Peptide having SEQ. ID. NO. 25

The above-identified peptide is assembled on 4-(2'-  
10 4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 5. Used in analysis are Solvent A  
15 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
20 calculated 4172.6.

#### EXAMPLE 22

##### Preparation of Peptide having SEQ. ID. NO. 26

25 The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a

similar way to Example 5. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4115.5.

EXAMPLE 23

Preparation of Peptide having SEQ. ID. NO. 27

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4188.6.

EXAMPLE 24Preparation of Peptide having SEQ. ID. NO. 28

The above-identified peptide is assembled on 4-(2'-  
5 4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 5. Used in analysis are Solvent A  
10 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
15 calculated 4131.6.

EXAMPLE 25Preparation of Peptide having SEQ. ID. NO. 29

The above-identified peptide is assembled on 4-(2'-  
20 4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
25 similar way to Example 5. Used in analysis are Solvent A  
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the



product peptide. Electrospray Mass Spectrometry (M):  
calculated 4172.6.

#### EXAMPLE 26

##### 5      Preparation of Peptide having SEQ. ID. NO. 30

The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
10      Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 5. Used in analysis are Solvent A  
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
15      Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 4145.6.

20

#### EXAMPLE 27

##### Preparation of Peptide having SEQ. ID. NO. 31

The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
25      norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 5. Additional double couplings are  
required at the thioproline positions 38, 37, 36 and 31.

Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4266.8.

#### EXAMPLE 28

##### Preparation of Peptide having SEQ. ID. NO. 32

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Additional double couplings are required at the thioproline positions 38, 37 and 36. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4246.8.

#### EXAMPLE 29

##### Preparation of Peptide having SEQ. ID. NO. 33

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide

norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Additional double couplings are required at the homoproline positions 38, 37, 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4250.8.

#### EXAMPLE 30

##### Preparation of Peptide having SEQ. ID. NO. 34

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Additional double couplings are required at the homoproline positions 38, 37, and 36. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4234.8.

EXAMPLE 31Preparation of Peptide having SEQ. ID. NO. 35

5           The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
10       similar way to Example 5. Additional double couplings are  
required at the thioproline positions 38, 37, 36 and 31.  
Used in analysis are Solvent A (0.1% TFA in water) and  
Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient  
30% to 60% Solvent B in Solvent A over 30 minutes) of the  
15       lyophilized peptide is then carried out to determine the  
retention time of the product peptide. Electrospray Mass  
Spectrometry (M): calculated 4209.8.

EXAMPLE 32Preparation of Peptide having SEQ. ID. NO. 36

20           The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
25       Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 5. Additional double couplings are  
required at the homoproline positions 38, 37, 36 and 31.  
Used in analysis are Solvent A (0.1% TFA in water) and

Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4193.7.

#### EXAMPLE 33

##### Preparation of Peptide having SEQ. ID. NO. 37

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Additional double couplings are required at the N-methylalanine positions 38, 37, 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3858.2.

#### EXAMPLE 34

##### Preparation of Peptide having SEQ. ID. NO. 38

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using

Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 5. Additional double couplings are  
required at the N-methylalanine positions 38, 37 and 36.

5 Used in analysis are Solvent A (0.1% TFA in water) and  
Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient  
30% to 60% Solvent B in Solvent A over 30 minutes) of the  
lyophilized peptide is then carried out to determine the  
retention time of the product peptide. Electrospray Mass  
10 Spectrometry (M): calculated 3940.3.

#### EXAMPLE 35

##### Preparation of Peptide having SEQ. ID. NO. 39

15 The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
20 cleaved from the resin, deprotected and purified in a  
similar way to Example 5. Additional double couplings are  
required at the N-methylalanine positions 38, 37, 36 and  
31. Used in analysis are Solvent A (0.1% TFA in water) and  
Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient  
25 30% to 60% Solvent B in Solvent A over 30 minutes) of the  
lyophilized peptide is then carried out to determine the  
retention time of the product peptide. Electrospray Mass  
Spectrometry (M): calculated 3801.1.

EXAMPLE 36Preparation of C-terminal carboxylic acid Peptides  
corresponding to the above C-terminal amide sequences.

5

The above peptides of Examples 5 to 35 are assembled on the so called Wang resin (p-alkoxybenzylalcohol resin (Bachem, 0.54 mmole/g)) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin,  
10 deprotected and purified in a similar way to Example 5. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the  
15 retention time of the product peptide. Electrospray Mass Spectrometry provides an experimentally determined (M).

EXAMPLE 37

20

Preparation of Peptide having SEQ ID NO. 7

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly-NH<sub>2</sub> [SEQ. ID. NO. 7]

25

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.). In  
30 general, single-coupling cycles were used throughout the

synthesis and Fast Moc (HBTU activation) chemistry was employed. Deprotection (Fmoc group removal) of the growing peptide chain was achieved using piperidine. Final deprotection of the completed peptide resin was achieved using a mixture of triethylsilane (0.2 mL), ethanedithiol (0.2 mL), anisole (0.2 mL), water (0.2 mL) and trifluoroacetic acid (15 mL) according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems, Inc.) The peptide was precipitated in ether/water (50 mL) and centrifuged. The precipitate was reconstituted in glacial acetic acid and lyophilized. The lyophilized peptide was dissolved in water). Crude purity was about 75%.

Used in purification steps and analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). The solution containing peptide was applied to a preparative C-18 column and purified (10% to 40% Solvent B in Solvent A over 40 minutes). Purity of fractions was determined isocratically using a C-18 analytical column. Pure fractions were pooled furnishing the above-identified peptide. Analytical RP-HPLC (gradient 30% to 50% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 18.9 minutes. Electrospray Mass Spectrometry (M): calculated 3408.0; found 3408.9.



EXAMPLE 38Preparation of Peptide having SEQ ID NO. 40

5 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 40]

The above amidated peptide was assembled on 4-(2'-4'-  
10 dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 37. Used in analysis were Solvent A  
15 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 40% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide gave  
product peptide having an observed retention time of 17.9  
minutes. Electrospray Mass Spectrometry (M): calculated  
20 3294.7; found 3294.8.

EXAMPLE 39Preparation of Peptide having SEQ ID NO. 41

25

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 41]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 29% to 36% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 20.7 minutes. Electrospray Mass Spectrometry (M): calculated 3237.6; found 3240.

#### EXAMPLE 40

##### Preparation of Peptide having SEQ ID NO. 42

His Ala Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 42]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).

Analytical RP-HPLC (gradient 36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 15.2 minutes. Electrospray Mass Spectrometry (M): calculated 3251.6; found 3251.5.

#### EXAMPLE 41

##### Preparation of Peptide having SEQ ID NO. 43

His Gly Glu Gly Ala Phe Thr Ser Asp Leu Ser Lys Gln  
Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys  
Asn-NH<sub>2</sub> [SEQ. ID. NO. 43]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 13.1 minutes. Electrospray Mass Spectrometry (M): calculated 3207.6; found 3208.3.

EXAMPLE 42Preparation of Peptide having SEQ ID NO. 44

His Gly Glu Gly Thr Ala Thr Ser Asp Leu Ser Lys Gln Leu  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 44]

The above amidated peptide was assembled on 4-(2'-4'-  
dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
10 norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 37. Used in analysis were Solvent  
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
15 Analytical RP-HPLC (gradient 35% to 45% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide gave  
product peptide having an observed retention time of 12.8  
minutes. Electrospray Mass Spectrometry (M): calculated  
3161.5; found 3163.

20

EXAMPLE 43Preparation of Peptide having SEQ ID NO. 45

His Gly Glu Gly Thr Phe Thr Ala Asp Leu Ser Lys Gln Leu  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 45]

The above-identified amidated peptide was assembled

on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
5 purified in a similar way to Example 37. Used in analysis  
were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 36% to 46% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
gave product peptide having an observed retention time of  
10 15.2 minutes. Electrospray Mass Spectrometry (M):  
calculated 3221.6; found 3222.7.

#### EXAMPLE 44

##### Preparation of Peptide having SEQ ID NO. 46

15

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 46]

20

The above-identified amidated peptide was assembled  
on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
25 purified in a similar way to Example 37. Used in analysis  
were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 34% to 44% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide

gave product peptide having an observed retention time of 14.3 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3199.4.

5

EXAMPLE 45Preparation of Peptide having SEQ ID NO. 47

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ala Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 47]

10

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 37. Used in analysis were Solvent A  
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 38% to 48% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide gave  
product peptide having an observed retention time of 15.7  
minutes. Electrospray Mass Spectrometry (M): calculated  
3221.6; found 3221.6.

15

20

25

EXAMPLE 46Preparation of Peptide having SEQ ID NO. 48

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Ala Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 48]

30

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 18.1 minutes. Electrospray Mass Spectrometry (M): calculated 3180.5; found 3180.9.

#### EXAMPLE 47

##### Preparation of Peptide having SEQ ID NO. 49

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Ala Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 49]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide

gave product peptide having an observed retention time of 17.0 minutes. Electrospray Mass Spectrometry (M): calculated 3180.6; found 3182.8.

5

EXAMPLE 48Preparation of Peptide having SEQ ID NO. 50

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Ala  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 50]

10

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 32% to 42% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.9 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3195.9.

15

20

25

EXAMPLE 49Preparation of Peptide having SEQ ID NO. 51

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu



Ala Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 51]

5 The above-identified amidated peptide was assembled  
on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
10 purified in a similar way to Example 37. Used in analysis  
were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
gave product peptide having an observed retention time of  
17.9 minutes. Electrospray Mass Spectrometry (M):  
15 calculated 3179.6; found 3179.0.

#### EXAMPLE 50

##### Preparation of Peptide having SEQ ID NO. 52

20

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Ala Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 52]

25

The above-identified amidated peptide was assembled  
on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and

purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.3 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3180.0.

#### EXAMPLE 51

##### Preparation of Peptide having SEQ ID NO. 53

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Ala Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 53]

The above-identified peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 13.7 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3179.0.

EXAMPLE 52Preparation of Peptide having SEQ ID NO. 54

5 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Ala Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 54]

The above-identified amidated peptide was assembled  
10 on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 37. Used in analysis  
15 were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 35% to 45% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
gave product peptide having an observed retention time of  
14.0 minutes. Electrospray Mass Spectrometry (M):  
20 calculated 3209.6; found 3212.8.

EXAMPLE 53Preparation of Peptide having SEQ ID NO. 55

25 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Ala Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 55]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.3 minutes. Electrospray Mass Spectrometry (M): calculated 3152.5; found 3153.5.

#### EXAMPLE 54

##### Preparation of Peptide having SEQ ID NO. 56

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Ala Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 56]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 35% to 45% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide

gave product peptide having an observed retention time of 12.1 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3197.7.

5

EXAMPLE 55Preparation of Peptide having SEQ ID NO. 57

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Ala Phe Leu Lys Asn-NH<sub>2</sub>  
10 [SEQ. ID. NO. 57]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55  
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B  
20 in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 10.9 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3180.5.

25

EXAMPLE 56Preparation of Peptide having SEQ ID NO. 58

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Ala Leu Lys Asn-NH<sub>2</sub>

[SEQ. ID. NO. 58]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 32% to 42% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 17.5 minutes. Electrospray Mass Spectrometry (M): calculated 3161.5; found 3163.0.

#### EXAMPLE 57

##### Preparation of Peptide having SEQ ID NO. 59

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Ala Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 59]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 32% to 42% Solvent B

in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 19.5 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3199.

5

#### EXAMPLE 58

##### Preparation of Peptide having SEQ ID NO. 60

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
10 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Ala Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 60]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
15 acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis  
20 were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.5 minutes. Electrospray Mass Spectrometry (M): calculated 3180.5; found 3183.7.

25

EXAMPLE 59Preparation of Peptide having SEQ ID NO. 61

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Ala-NH<sub>2</sub>  
[SEQ. ID. NO. 61]

The above-identified amidated peptide was assembled  
on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 37. Used in analysis  
15 were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 34% to 44% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
gave product peptide having an observed retention time of  
22.8 minutes. Electrospray Mass Spectrometry (M):  
20 calculated 3194.6; found 3197.6.

EXAMPLE 60Preparation of Peptide having SEQ ID NO. 62

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro-NH<sub>2</sub> [SEQ. ID. NO.  
62]



The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4099.6.

#### EXAMPLE 61

##### Preparation of Peptide having SEQ ID NO. 63

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro-NH<sub>2</sub> [SEQ. ID. NO.  
63]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA

in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):  
5 calculated 4042.5.

#### EXAMPLE 62

##### Preparation of Peptide having SEQ ID NO. 64

10 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala Pro Pro-NH<sub>2</sub> [SEQ. ID. NO. 64]

The above-identified peptide is assembled on 4-(2'-  
15 4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 37. Used in analysis are Solvent A  
20 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
25 calculated 4002.4

EXAMPLE 63Preparation of Peptide having SEQ ID NO. 65

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala Pro Pro-NH<sub>2</sub> [SEQ. ID. NO. 65]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 37. Used in analysis  
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
20 calculated 3945.4.

EXAMPLE 64Preparation of Peptide having SEQ ID NO. 66

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln  
25 Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys  
Asn Gly Gly Pro Ser Ser Gly Ala Pro-NH<sub>2</sub> [SEQ. ID. NO. 66]

The above-identified amidated peptide is assembled on

4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
5 purified in a similar way to Example 37. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
10 product peptide. Electrospray Mass Spectrometry (M):  
calculated 3905.3.

#### EXAMPLE 65

##### Preparation of Peptide having SEQ ID NO. 67

15 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala Pro-NH<sub>2</sub> [SEQ. ID. NO. 67]

20 The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
25 purified in a similar way to Example 37. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3848.2.

5

EXAMPLE 66Preparation of Peptide having SEQ ID NO. 68

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
10 Gly Gly Pro Ser Ser Gly Ala-NH<sub>2</sub> [SEQ. ID. NO. 68]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55  
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3808.2.

25

EXAMPLE 67Preparation of Peptide having SEQ ID NO. 69

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala-NH<sub>2</sub> [SEQ. ID. NO. 69]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 37. Used in analysis  
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3751.1.

20

EXAMPLE 68Preparation of Peptide having SEQ ID NO. 70

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly-NH<sub>2</sub> [SEQ. ID. NO. 70]

The above-identified amidated peptide is assembled on

4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
5 purified in a similar way to Example 37. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
10 product peptide. Electrospray Mass Spectrometry (M):  
calculated 3737.1.

#### EXAMPLE 69

##### Preparation of Peptide having SEQ ID NO. 71

15 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro Ser Ser Gly-NH<sub>2</sub> [SEQ. ID. NO. 71]

20 The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
25 purified in a similar way to Example 37. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3680.1.

5

EXAMPLE 70Preparation of Peptide having SEQ ID NO. 72

10

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser-NH<sub>2</sub> [SEQ. ID. NO. 72]

15

20

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3680.1

25

EXAMPLE 71Preparation of Peptide having SEQ ID NO. 73

30

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro Ser Ser-NH<sub>2</sub> [SEQ. ID. NO. 73]



The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3623.0.

#### EXAMPLE 72

##### Preparation of Peptide having SEQ ID NO. 74

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser-NH<sub>2</sub> [SEQ. ID. NO. 74]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA

in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):  
5 calculated 3593.0

### EXAMPLE 73

#### Preparation of Peptide having SEQ ID NO. 75

10 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro Ser-NH<sub>2</sub> [SEQ. ID. NO. 75]

The above-identified amidated peptide is assembled on  
15 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 37. Used in analysis  
20 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
25 calculated 3535.9

EXAMPLE 74Preparation of Peptide having SEQ ID NO. 76

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro-NH<sub>2</sub> [SEQ. ID. NO. 76]

The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
10 norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 37. Used in analysis are Solvent A  
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
15 Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3505.9.

20

EXAMPLE 75Preparation of Peptide having SEQ ID NO. 77

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro-NH<sub>2</sub> [SEQ. ID. NO. 77]

The above-identified amidated peptide is assembled on

4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
5 purified in a similar way to Example 37. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
10 product peptide. Electrospray Mass Spectrometry (M):  
calculated 3448.8.

#### EXAMPLE 76

##### Preparation of Peptide having SEQ ID NO. 78

15 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly-NH<sub>2</sub> [SEQ. ID. NO. 78]

20 The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
25 similar way to Example 37. Used in analysis are Solvent A  
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is

then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3351.7.

5

EXAMPLE 77Preparation of Peptide having SEQ ID NO. 79

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
10 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly-NH<sub>2</sub> [SEQ. ID. NO. 79]

The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
15 norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 37. Used in analysis are Solvent A  
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
20 Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3351.8.

25

EXAMPLE 78Preparation of Peptide having SEQ ID NO. 80

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly-NH<sub>2</sub> [SEQ. ID. NO. 80]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 37. Used in analysis  
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
20 calculated 3294.7.

EXAMPLE 79Preparation of Peptide having SEQ ID NO. 81

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly tPro Ser Ser Gly Ala tPro tPro tPro-NH<sub>2</sub> [SEQ. ID.  
NO. 81]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Double couplings are required at residues 37, 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4197.1.

15

EXAMPLE 80Preparation of Peptide having SEQ ID NO. 82

20

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala tPro tPro tPro-NH<sub>2</sub> [SEQ. ID.  
NO. 82]

25

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Double couplings

are required at residues 37, 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4179.1.

#### EXAMPLE 81

##### Preparation of Peptide having SEQ ID NO. 83

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly NMeala Ser Ser Gly Ala Pro Pro-NH<sub>2</sub> [SEQ. ID. NO.  
83]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Double couplings are required at residues 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):



calculated 3948.3.

EXAMPLE 82

Preparation of Peptide having SEQ ID NO. 84

5

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly NMeala Ser Ser Gly Ala NMeala Nmeala-NH<sub>2</sub> [SEQ. ID.  
NO. 84]

10

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
15 Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 37. Double couplings  
are required at residues 36 and 31. Used in analysis are  
Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in  
ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
20 Solvent A over 30 minutes) of the lyophilized peptide is  
then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3840.1.

EXAMPLE 83Preparation of Peptide having SEQ ID NO. 85

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly hPro Ser Ser Gly Ala hPro hPro-NH<sub>2</sub> [SEQ. ID. NO.  
85]

The above-identified amidated peptide is assembled on  
10 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
15 purified in a similar way to Example 37. Double couplings  
are required at residues 36 and 31. Used in analysis are  
Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in  
ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide is  
20 then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 4050.1.

EXAMPLE 84Preparation of Peptide having SEQ ID NO. 86

25

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly hPro Ser Ser Gly Ala hPro-NH<sub>2</sub> [SEQ. ID. NO. 86]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. A double coupling is required at residue 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3937.1

#### EXAMPLE 85

##### Preparation of Peptide having SEQ ID NO. 87

Arg Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala-NH<sub>2</sub> [SEQ. ID. NO. 87]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis

are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3827.2.

#### EXAMPLE 86

##### Preparation of Peptide having SEQ ID NO. 88

His Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly-NH<sub>2</sub> [SEQ. ID. NO. 88]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3394.8.

EXAMPLE 87Preparation of Peptide having SEQ ID NO. 89

His Gly Glu Gly Thr Naphthylala Thr Ser Asp Leu Ser Lys  
5 Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu  
Lys Asn-NH<sub>2</sub> [SEQ. ID. NO. 89]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 37. Used in analysis  
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
20 calculated 3289.5.

EXAMPLE 88Preparation of Peptide having SEQ ID NO. 90

His Gly Glu Gly Thr Phe Ser Ser Asp Leu Ser Lys Gln Met  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 90]

The above-identified amidated peptide is assembled on

4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
5 purified in a similar way to Example 37. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
10 product peptide. Electrospray Mass Spectrometry (M):  
calculated 3280.7.

#### EXAMPLE 89

##### Preparation of Peptide having SEQ ID NO. 91

15 His Gly Glu Gly Thr Phe Ser Thr Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 91]

20 The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
25 purified in a similar way to Example 37. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3294.7.

5

EXAMPLE 90Preparation of Peptide having SEQ ID NO. 92

His Gly Glu Gly Thr Phe Thr Ser Glu Leu Ser Lys Gln Met  
Ala Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
10 [SEQ. ID. NO. 92]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and  
15 purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide  
20 is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3250.7.

EXAMPLE 91

25

Preparation of Peptide having SEQ ID NO. 93

His Gly Glu Gly Thr Phe Thr Ser Asp pentylgly Ser Lys Gln  
Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys

Asn-NH<sub>2</sub> [SEQ. ID. NO. 93]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3253.5.

#### EXAMPLE 92

##### Preparation of Peptide having SEQ ID NO. 94

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Naphthylala Ile Glu Phe Leu  
Lys Asn-NH<sub>2</sub> [SEQ. ID. NO. 94]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis



are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3289.5.

### EXAMPLE 93

#### Preparation of Peptide having SEQ ID NO. 95

10

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe tButylgly Glu Trp Leu Lys  
Asn-NH<sub>2</sub> [SEQ. ID. NO. 95]

15

20

25

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3183.4.

EXAMPLE 94Preparation of Peptide having SEQ ID NO. 96

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Asp Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 96]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 37. Used in analysis  
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
20 calculated 3237.6.

EXAMPLE 95Preparation of Peptide having SEQ ID NO. 97

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Leu  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro Ser Ser-NH<sub>2</sub> [SEQ. ID. NO. 97]

The above-identified amidated peptide is assembled on

4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
5 purified in a similar way to Example 37. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
10 product peptide. Electrospray Mass Spectrometry (M):  
calculated 3637.9.

#### EXAMPLE 96

##### Preparation of Peptide having SEQ ID NO. 98

15 His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly-NH<sub>2</sub> [SEQ. ID. NO. 98]

20 The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
25 purified in a similar way to Example 37. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3309.7.

EXAMPLE 97

Preparation of Peptide having SEQ ID NO. 99

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly hPro Ser Ser Gly Ala hPro hPro-NH<sub>2</sub> [SEQ. ID. NO.  
99]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Double couplings are required at residues 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3711.1.

EXAMPLE 98

Preparation of C-terminal carboxylic acid peptides corresponding to the above C-terminal amide sequences for SEQ ID NOS. 7, 40-61, 68-75, 78-80 and 87-96

5

Peptides having the sequences of SEQ ID NOS. 7, 40-61, 68-75, 78-80 and 87-96 are assembled on the so called Wang resin (p-alkoxybenzylalcohol resin (Bachem, 0.54 mmole/g)) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry provides an experimentally determined (M).

10

15

20

EXAMPLE 99

Preparation of C-terminal carboxylic acid peptides corresponding to the above C-terminal amide sequences for SEQ ID NOS. 62-67, 76, 77 and 81-86

25

Peptides having the sequences of SEQ ID NOS. 62-67, 76, 77 and 81-86 are assembled on the 2-chlorotritylchloride resin (200-400 mesh), 2% DVB (Novabiochem, 0.4-1.0 mmole/g)) using Fmoc-protected amino

acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry provides an experimentally determined (M).

#### EXAMPLE 100

##### Preparation of Peptide having SEQ ID NO. 100

Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 100]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.). In general, single-coupling cycles were used throughout the synthesis and Fast Moc (HBTU activation) chemistry was employed. Deprotection (Fmoc group removal) of the growing peptide chain was achieved using piperidine. Final deprotection of the completed peptide resin was achieved using a mixture of triethylsilane (0.2 mL), ethanedithiol (0.2 mL), anisole (0.2 mL), water (0.2 mL) and trifluoroacetic acid (15 mL) according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems,

Inc.) The peptide was precipitated in ether/water (50 mL) and centrifuged. The precipitate was reconstituted in glacial acetic acid and lyophilized. The lyophilized peptide was dissolved in water). Crude purity was about 75%.

Used in purification steps and analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).

The solution containing peptide was applied to a preparative C-18 column and purified (10% to 40% Solvent B in Solvent A over 40 minutes). Purity of fractions was determined isocratically using a C-18 analytical column. Pure fractions were pooled furnishing the above-identified peptide. Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 19.2 minutes. Electrospray Mass Spectrometry (M): calculated 3171.6; found 3172.

#### EXAMPLE 101

##### Preparation of Peptide having SEQ ID NO. 101

His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 101]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.),

cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.9 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3180.

10

EXAMPLE 102Preparation of Peptide having SEQ ID NO. 102

His Gly Glu Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 102]

15

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 12.2 minutes. Electrospray Mass Spectrometry (M): calculated 3251.6; found 3253.3.

20

25



EXAMPLE 103Preparation of Peptide having SEQ ID NO. 103

5 His Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 103]

The above amidated peptide was assembled on 4-(2'-4'-  
10 dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide  
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using  
Fmoc-protected amino acids (Applied Biosystems, Inc.),  
cleaved from the resin, deprotected and purified in a  
similar way to Example 100. Used in analysis were Solvent  
15 A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).  
Analytical RP-HPLC (gradient 35% to 45% Solvent B in  
Solvent A over 30 minutes) of the lyophilized peptide gave  
product peptide having an observed retention time of 16.3  
minutes. Electrospray Mass Spectrometry (M): calculated  
20 3193.6; found 3197.

EXAMPLE 104Preparation of Peptide having SEQ ID NO. 104

25 Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 104]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3228.6.

#### EXAMPLE 105

##### Preparation of Peptide having SEQ ID NO. 105

His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 105]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3234.7.

5

EXAMPLE 106Preparation of Peptide having SEQ ID NO. 106

His Gly Glu Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 106]

10

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3308.7.

15

20

25

EXAMPLE 107Preparation of Peptide having SEQ ID NO. 107

His Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>

[SEQ. ID. NO. 107]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3250.7

#### EXAMPLE 108

##### Preparation of Peptide having SEQ ID NO. 108

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 108]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA

in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):  
5 calculated 3252.6.

#### EXAMPLE 109

##### Preparation of Peptide having SEQ ID NO. 109

10 Ala Ala Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 109]

The above-identified amidated peptide is assembled on  
15 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
20 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
25 calculated 3200.6.

EXAMPLE 110Preparation of Peptide having SEQ ID NO. 110

Ala Ala Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
5 [SEQ. ID. NO. 110]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
10 mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
15 in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3143.5.

20

EXAMPLE 111Preparation of Peptide having SEQ ID NO. 111

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 111]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3214.6.

#### EXAMPLE 112

##### Preparation of Peptide having SEQ ID NO. 112

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub> [SEQ. ID. NO. 112]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):

calculated 3157.5.

EXAMPLE 113

Preparation of Peptide having SEQ ID NO. 113

5

Ala Gly Asp Gly Ala Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 113]

10

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
15 purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
20 product peptide. Electrospray Mass Spectrometry (M):  
calculated 3184.6.

EXAMPLE 114

Preparation of Peptide having SEQ ID NO. 114

25

Ala Gly Asp Gly Ala Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 114]



The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3127.5.

#### EXAMPLE 115

##### Preparation of Peptide having SEQ ID NO. 115

Ala Gly Asp Gly Thr NaphthylAla Thr Ser Asp Leu Ser Lys  
Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu  
Lys Asn-NH<sub>2</sub> [SEQ. ID. NO. 115]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B

in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3266.4.

5

EXAMPLE 116Preparation of Peptide having SEQ ID NO. 116

Ala Gly Asp Gly Thr Naphthylala Thr Ser Asp Leu Ser Lys  
10 Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu  
Lys Asn-NH<sub>2</sub> [SEQ. ID. NO. 116]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
15 acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3209.4.

25

EXAMPLE 117Preparation of Peptide having SEQ ID NO. 117

Ala Gly Asp Gly Thr Phe Ser Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 117]

5           The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
10       purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
15       product peptide. Electrospray Mass Spectrometry (M):  
calculated 3200.6.

#### EXAMPLE 118

##### Preparation of Peptide having SEQ ID NO. 118

20

Ala Gly Asp Gly Thr Phe Ser Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 118]

25

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and

purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
5 is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3143.5.

#### EXAMPLE 119

#### 10 Preparation of Peptide having SEQ ID NO. 119

Ala Gly Asp Gly Thr Phe Thr Ala Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 119]

15 The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
20 Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
25 is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3198.6.

EXAMPLE 120Preparation of Peptide having SEQ ID NO. 120

Ala Gly Asp Gly Thr Phe Thr Ala Asp Leu Ser Lys Gln Leu  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 120]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
15 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3141.5.

20

EXAMPLE 121Preparation of Peptide having SEQ ID NO. 121

Ala Gly Asp Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Met  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 121]

The above-identified peptide is assembled on 4-(2'-  
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide

norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3170.6.

#### EXAMPLE 122

##### Preparation of Peptide having SEQ ID NO. 122

Ala Gly Asp Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 122]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):

calculated 3113.5.

EXAMPLE 123

Preparation of Peptide having SEQ ID NO. 123

5

Ala Gly Asp Gly Thr Phe Thr Ser Glu Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 123]

10

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
15 purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
20 product peptide. Electrospray Mass Spectrometry (M):  
calculated 3228.6.

EXAMPLE 124

Preparation of Peptide having SEQ ID NO. 124

25

Ala Gly Asp Gly Thr Phe Thr Ser Glu Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 124]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3171.6.

#### EXAMPLE 125

##### Preparation of Peptide having SEQ ID NO. 125

Ala Gly Asp Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 125]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B



in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3172.5.

5

EXAMPLE 126Preparation of Peptide having SEQ ID NO. 126

Ala Gly Asp Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Leu  
10 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 126]

The above-identified amidated peptiden is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
15 acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3115.4.

25

EXAMPLE 127Preparation of Peptide having SEQ ID NO. 127

Ala Gly Asp Gly Thr Phe Thr Ser Asp Pentylgly Ser Lys Gln  
5 Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys  
Asn-NH<sub>2</sub> [SEQ. ID. NO. 127]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3230.4.

EXAMPLE 128Preparation of Peptide having SEQ ID NO. 128

Ala Gly Asp Gly Thr Phe Thr Ser Asp Pentylgly Ser Lys Gln  
25 Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys  
Asn-NH<sub>2</sub> [SEQ. ID. NO. 128]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3198.6.

#### EXAMPLE 129

##### Preparation of Peptide having SEQ ID NO. 129

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ala Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 129]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):

calculated 3141.5.

EXAMPLE 130

Preparation of Peptide having SEQ ID NO. 130

5

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ala Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 130]

10

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
15 purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
20 product peptide. Electrospray Mass Spectrometry (M):  
calculated 3157.5.

EXAMPLE 131

Preparation of Peptide having SEQ ID NO. 131

25

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Ala Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 131]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3100.4.

#### EXAMPLE 132

##### Preparation of Peptide having SEQ ID NO. 132

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Ala Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 132]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B

in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3157.6.

5

### EXAMPLE 133

#### Preparation of Peptide having SEQ ID NO. 133

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Ala Met  
10 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 133]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
15 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3100.5.

25

### EXAMPLE 134

#### Preparation of Peptide having SEQ ID NO. 134

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Ala Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 134]

5           The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
10       purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
15       product peptide. Electrospray Mass Spectrometry (M):  
calculated 3100.5.

#### EXAMPLE 135

##### Preparation of Peptide having SEQ ID NO. 135

20

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Ala  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 135]

25

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and

purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3154.5.

#### EXAMPLE 136

##### Preparation of Peptide having SEQ ID NO. 136

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Ala  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 136]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3115.5.



EXAMPLE 137Preparation of Peptide having SEQ ID NO. 137

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln  
5 Pentylgly Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu  
Lys Asn-NH<sub>2</sub> [SEQ. ID. NO. 137]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
20 calculated 3212.4.

EXAMPLE 138Preparation of Peptide having SEQ ID NO. 138

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln  
25 Pentylgly Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu  
Lys Asn-NH<sub>2</sub> [SEQ. ID. NO. 138]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3173.4.

#### EXAMPLE 139

##### Preparation of Peptide having SEQ ID NO. 139

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Ala Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 139]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3156.6.

EXAMPLE 140Preparation of Peptide having SEQ ID NO. 140

5     Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Ala Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 140]

10     The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
15     are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
20     calculated 3099.5.

EXAMPLE 141Preparation of Peptide having SEQ ID NO. 141

25     Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Ala Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 141]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3156.6.

#### EXAMPLE 142

##### Preparation of Peptide having SEQ ID NO. 142

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Ala Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 142]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the

product peptide. Electrospray Mass Spectrometry (M):  
calculated 3099.5.

EXAMPLE 143

5           Preparation of Peptide having SEQ ID NO. 143

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Ala Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 143]

10

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
15 Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
20 is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3156.6.

EXAMPLE 144

25           Preparation of Peptide having SEQ ID NO. 144

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Ala Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 144]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3099.5.

#### EXAMPLE 145

##### Preparation of Peptide having SEQ ID NO. 145

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Ala Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 145]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3186.6.

5

EXAMPLE 146Preparation of Peptide having SEQ ID NO. 146

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Ala Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
10 [SEQ. ID. NO. 146]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55  
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
20 in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3129.5.

25

EXAMPLE 147Preparation of Peptide having SEQ ID NO. 147

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Ala Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 147]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3129.5.

15

EXAMPLE 148Preparation of Peptide having SEQ ID NO. 148

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Ala Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 148]

20

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B

25



in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3072.4.

5

EXAMPLE 149

Preparation of Peptide having SEQ ID NO. 149

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
10 Glu Glu Glu Ala Val Arg Ala Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 149]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
15 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3172.5.

25

EXAMPLE 150Preparation of Peptide having SEQ ID NO. 150

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
5 Glu Glu Glu Ala Val Arg Ala Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 150]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
20 calculated 3115.5.

EXAMPLE 151Preparation of Peptide having SEQ ID NO. 151

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
25 Glu Glu Glu Ala Val Arg Leu Naphthylala Ile Glu Trp Leu  
Lys Asn-NH<sub>2</sub> [SEQ. ID. NO. 151]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis  
5 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):  
10 calculated 3266.4.

#### EXAMPLE 152

##### Preparation of Peptide having SEQ ID NO. 152

15 Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Naphthylala Ile Glu Phe Leu  
Lys Asn-NH<sub>2</sub> [SEQ. ID. NO. 152]

The above-identified amidated peptide is assembled on  
20 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis  
25 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):

calculated 3209.4.

EXAMPLE 153

Preparation of Peptide having SEQ ID NO. 153

5

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Val Glu Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 153]

10

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
15 purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
20 product peptide. Electrospray Mass Spectrometry (M):  
calculated 3200.6.

EXAMPLE 154

Preparation of Peptide having SEQ ID NO. 154

25

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Val Glu Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 154]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3143.5.

#### EXAMPLE 155

##### Preparation of Peptide having SEQ ID NO. 155

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe tButylgly Glu Trp Leu Lys  
Asn-NH<sub>2</sub> [SEQ. ID. NO. 155]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B

in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3216.5.

5

EXAMPLE 156Preparation of Peptide having SEQ ID NO. 156

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
10 Glu Glu Glu Ala Val Arg Leu Phe tButylgly Glu Phe Leu Lys  
Asn-NH<sub>2</sub> [SEQ. ID. NO. 156]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
15 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3159.4.

25

EXAMPLE 157Preparation of Peptide having SEQ ID NO. 157

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Asp Trp Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 157]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
15 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3200.6.

20

EXAMPLE 158Preparation of Peptide having SEQ ID NO. 158

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Asp Phe Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 158]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3143.5.

#### EXAMPLE 159

##### Preparation of Peptide having SEQ ID NO. 159

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Ala Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 159]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):



calculated 3099.5.

EXAMPLE 160

Preparation of Peptide having SEQ ID NO. 160

5

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Ala Leu Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 160]

10

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
15 purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
20 product peptide. Electrospray Mass Spectrometry (M):  
calculated 3081.4.

EXAMPLE 161

Preparation of Peptide having SEQ ID NO. 161

25

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Ala Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 161]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3172.5.

#### EXAMPLE 162

##### Preparation of Peptide having SEQ ID NO. 162

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Ala Lys Asn-NH<sub>2</sub>  
[SEQ. ID. NO. 162]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3115.5.

5

EXAMPLE 163Preparation of Peptide having SEQ ID NO. 163

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Ala Asn-NH<sub>2</sub>  
10 [SEQ. ID. NO. 163]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55  
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
20 in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3157.5.

25

EXAMPLE 164Preparation of Peptide having SEQ ID NO. 164

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Ala Asn-NH<sub>2</sub>

[SEQ. ID. NO. 164]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
5 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
10 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3100.4.

15

EXAMPLE 165

Preparation of Peptide having SEQ ID NO. 165

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
20 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Ala-NH<sub>2</sub>  
[SEQ. ID. NO. 165]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
25 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA

in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):  
5 calculated 3171.6.

#### EXAMPLE 166

##### Preparation of Peptide having SEQ ID NO. 166

10 Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Ala-NH<sub>2</sub>  
[SEQ. ID. NO. 166]

The above-identified amidated peptide is assembled on  
15 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
20 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
25 calculated 3114.5.

EXAMPLE 167Preparation of Peptide having SEQ ID NO. 167

Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro-NH<sub>2</sub> [SEQ. ID. NO.  
167]

The above-identified amidated peptide is assembled on  
10 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
20 calculated 4033.5.

EXAMPLE 168Preparation of Peptide having SEQ ID NO. 168

His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro-NH<sub>2</sub> [SEQ. ID. NO.  
168]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3984.4.

#### EXAMPLE 169

##### Preparation of Peptide having SEQ ID NO. 169

His Gly Glu Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala Pro Pro-NH<sub>2</sub> [SEQ. ID. NO. 169]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4016.5.

5

EXAMPLE 170Preparation of Peptide having SEQ ID NO. 170

His Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
10 Gly Gly Pro Ser Ser Gly Ala Pro-NH<sub>2</sub> [SEQ. ID. NO. 170]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55  
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
20 in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3861.3.

25

EXAMPLE 171Preparation of Peptide having SEQ ID NO. 171

Ala Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn



Gly Gly Pro Ser Ser Gly Ala Pro-NH<sub>2</sub> [SEQ. ID. NO. 171]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3746.1.

#### EXAMPLE 172

##### Preparation of Peptide having SEQ ID NO. 172

Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala-NH<sub>2</sub> [SEQ. ID. NO. 172]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA

in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):  
5 calculated 3742.1.

EXAMPLE 173

Preparation of Peptide having SEQ ID NO. 173

10 His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala-NH<sub>2</sub> [SEQ. ID. NO. 173]

The above-identified amidated peptide is assembled on  
15 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
20 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
25 calculated 3693.1.

EXAMPLE 174Preparation of Peptide having SEQ ID NO. 174

His Gly Glu Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly-NH<sub>2</sub> [SEQ. ID. NO. 174]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
20 calculated 3751.2.

EXAMPLE 175Preparation of Peptide having SEQ ID NO. 175

His Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Met  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser-NH<sub>2</sub> [SEQ. ID. NO. 175]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3634.1.

#### EXAMPLE 176

##### Preparation of Peptide having SEQ ID NO. 176

Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser-NH<sub>2</sub> [SEQ. ID. NO. 176]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. (Electrospray Mass Spectrometry (M):

calculated 3526.9.

EXAMPLE 177

Preparation of Peptide having SEQ ID NO. 177

5

His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro Ser-NH<sub>2</sub> [SEQ. ID. NO. 177]

10

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
15 purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
20 product peptide. Electrospray Mass Spectrometry (M):  
calculated 3477.9.

EXAMPLE 178

Preparation of Peptide having SEQ ID NO. 178

25

His Gly Glu Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro-NH<sub>2</sub> [SEQ. ID. NO. 178]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3519.9.

#### EXAMPLE 179

##### Preparation of Peptide having SEQ ID NO. 179

His Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly-NH<sub>2</sub> [SEQ. ID. NO. 179]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B

in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3307.7.

5

EXAMPLE 180Preparation of Peptide having SEQ ID NO. 180

Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
10 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly-NH<sub>2</sub> [SEQ. ID. NO. 180]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
15 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3186.5.

25

EXAMPLE 181Preparation of Peptide having SEQ ID NO. 181

His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly tPro Ser Ser Gly Ala tPro tPro tPro-NH<sub>2</sub> [SEQ. ID.  
NO. 181]

5

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
10 Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Double couplings  
are required at residues 37, 36 and 31. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
15 in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 4121.1.

20

#### EXAMPLE 182

##### Preparation of Peptide having SEQ. ID NO. 182

His Gly Glu Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala tPro tPro tPro-NH<sub>2</sub> [SEQ. ID.  
25 NO. 182].

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55



mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Double couplings are required at residues 37, 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4173.2.

#### EXAMPLE 183

##### Preparation of Peptide having SEQ ID NO. 183

His Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly NMeala Ser Ser Gly Ala NMeala NMeala-NH<sub>2</sub> [SEQ. ID.  
NO. 183]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Double couplings are required at residues 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is

then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3796.1.

5

EXAMPLE 184Preparation of Peptide having SEQ ID NO. 184

10

Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly hPro Ser Ser Gly Ala hPro-NH<sub>2</sub> [SEQ. ID. NO. 184]

15

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. A double coupling is required at residue 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3871.1.

25

EXAMPLE 185Preparation of Peptide having SEQ ID NO. 185

His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala-NH<sub>2</sub> [SEQ. ID. NO. 185]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
10 acetamide norleucine MBHA resin (Novabiochem, 0.55  
mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
15 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 3750.2.

20

EXAMPLE 186Preparation of Peptide having SEQ ID NO. 186

His Gly Asp Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly-NH<sub>2</sub> [SEQ. ID. NO. 186]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3408.8.

#### EXAMPLE 187

##### Preparation of Peptide having SEQ ID NO. 187

Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser-NH<sub>2</sub> [SEQ. ID.  
NO. 187]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the

product peptide. Electrospray Mass Spectrometry (M):  
calculated 4120.6.

#### EXAMPLE 188

##### 5           Preparation of Peptide having SEQ ID NO. 188

Ala Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu  
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn  
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser-NH<sub>2</sub> [SEQ. ID.  
10       NO. 188]

The above-identified amidated peptide is assembled on  
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy  
acetamide norleucine MBHA resin (Novabiochem, 0.55  
15       mmole/g) using Fmoc-protected amino acids (Applied  
Biosystems, Inc.), cleaved from the resin, deprotected and  
purified in a similar way to Example 100. Used in analysis  
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA  
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B  
20       in Solvent A over 30 minutes) of the lyophilized peptide  
is then carried out to determine the retention time of the  
product peptide. Electrospray Mass Spectrometry (M):  
calculated 4005.5.

25

#### EXAMPLE 189

Preparation of C-terminal carboxylic acid peptides  
corresponding to the above C-terminal amide sequences for  
Peptides having SEQ ID NOS. 100-166, 172-177, 179-180 and  
185-188.

C-terminal carboxylic acid peptides corresponding to amidated having SEQ ID NOS. 100-166, 172-177, 179-180 and 185-188 are assembled on the so called Wang resin (p-alkoxybenzylalcohol resin (Bachem, 0.54 mmole/g)) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to that described in Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry provides an experimentally determined (M).

#### EXAMPLE 190

Preparation of C-terminal carboxylic acid peptides corresponding to the above C-terminal amide sequences for Peptides having SEQ ID NOS. 167-171, 178 and 181-184.

C-terminal carboxylic acid peptides corresponding to amidated SEQ ID NOS. 167-171, 178 and 181-184 are assembled on the 2-chlorotritylchloride resin (200-400 mesh), 2% DVB (Novabiochem, 0.4-1.0 mmole/g)) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to that described in Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in

ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry provides  
5 an experimentally determined (M).

Various modifications of the invention in addition to those shown and described herein will become apparent to  
10 those skilled in the art from the foregoing description and fall within the scope of the following claims.

WE CLAIM:

1. A method for treating conditions or disorders which can be alleviated by reducing food intake in a subject comprising administering to said subject a  
5 therapeutically effective amount of an exendin or an exendin agonist.

2. The method according to claim 1 wherein said exendin or exendin agonist is administered parenterally.

3. The method according to claim 2 wherein said  
10 parenteral administration is by injection.

4. The method according to claim 3 wherein the injection is a peripheral injection.

5. The method according to claim 1 wherein about 10  
15  $\mu\text{g}$ -30 $\mu\text{g}$  to about 5mg of the exendin or exendin agonist is administered per day.

6. The method according to claim 1 wherein about 10  
 $\mu\text{g}$ -30  $\mu\text{g}$  to about 2 mg of the exendin or exendin agonist is administered per day.

7. The method according to claim 1, wherein about 30  
20  $\mu\text{g}$  to about 500  $\mu\text{g}$  of the exendin or exendin agonist is administered per day.

8. The method of claim 1 wherein said condition or disorder is obesity.

9. The method of claim 1 wherein said condition or  
25 disorder is Type II diabetes.

10. The method of claim 1 wherein said subject is human.

11. The method of claim 1 wherein said condition or disorder is an eating disorder.



12. The method of claim 1 wherein said condition or disorder is insulin-resistance syndrome.

13. A method for reducing the appetite of a subject comprising administering to said subject an appetite-lowering amount of an exendin or an exendin agonist.

14. A method for reducing the weight of a subject comprising administering to said subject a therapeutically effective amount of an exendin or an exendin agonist.

15. A method for lowering plasma lipids comprising administering to said subject a therapeutically effective amount of an exendin or an exendin agonist.

16. The method according to any of claims 1-15 wherein said exendin is exendin-3.

17. The method according to any of claims 1-15 wherein said exendin is exendin-4.

18. The method according to any of claims 1-15 wherein said exendin agonist is selected from the group consisting of exendin-4 (1-30), exendin-4 (1-30) amide, exendin-4 (1-28) amide, <sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 amide, <sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 (1-28) amide, and <sup>14</sup>Leu, <sup>22</sup>Ala, <sup>25</sup>Phe exendin-4 (1-28) amide.

19. The method according to any of claims 1-15, further comprising administering a therapeutically effective amount of one or more compounds selected from the group consisting essential of an amylin agonist, a leptin, and a CCK.

20. The method according to any of claims 1-15 wherein said exendin agonist is an exendin agonist according to Formula I.

21. The method according to any of claims 1-15 wherein said exendin agonist is an exendin agonist according to Formula II.

22. The method according to any of claims 1-15 wherein said exendin agonist is an exendin agonist according to Formula III.

23. A pharmaceutical composition for use in the treatment of conditions or disorders associated with hypernutrition comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.

24. The pharmaceutical composition according to claim 21, wherein said exendin is exendin-3.

25. The pharmaceutical composition according to claim 21 wherein said exendin is exendin-4.

26. The pharmaceutical composition according to claim 21 wherein said exendin agonist is selected from the group consisting of exendin-4 (1-30), exendin-4 (1-30) amide, exendin-4 (1-28) amide, <sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 amide, <sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 (1-28) amide, and <sup>14</sup>Leu, <sup>22</sup>Ala, <sup>25</sup>Phe exendin-4 (1-28) amide.

27. The pharmaceutical composition of claim 21 wherein said therapeutically effective amount is a therapeutically effective amount for a human subject.

28. A pharmaceutical composition for use in reducing the appetite of a subject comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.


29. A pharmaceutical composition for use in reducing the weight of a subject comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.

5        30. A pharmaceutical composition for use in lowering the plasma lipid level of a subject comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.

10       31. The pharmaceutical composition according to any of claims 21-28, further comprising a therapeutically effective amount of one or more compounds selected from the group consisting essentially of an amylin agonist, a leptin, and a CCK.


1/11

FIG. 1.






2/11

FIG. 2.




3/11

FIG. 3.




4/11

FIG. 4.



5/11


FIG. 5.






6/11

FIG. 6.




7/11

FIG. 7.




8/11

FIG. 8.



9/11

FIG. 9.



10/11

1 5 10 15 20  
 Xaa<sub>1</sub>, Xaa<sub>2</sub>, Xaa<sub>3</sub>, Gly Thr Xaa<sub>4</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>, Xaa<sub>7</sub>, Xaa<sub>8</sub>, Ser Lys Gln Xaa<sub>9</sub>, Glu Glu Glu Ala Val Arg Leu  
 30  
 Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Leu Lys Asn Gly Gly Xaa<sub>14</sub>, Ser Ser Gly Ala Xaa<sub>15</sub>, Xaa<sub>16</sub>, Xaa<sub>17</sub>, Xaa<sub>18</sub>-Z  
 35

(SEQ. ID. NO.)	Xaa <sub>1</sub>	Xaa <sub>2</sub>	Xaa <sub>3</sub>	Xaa <sub>4</sub>	Xaa <sub>5</sub>	Xaa <sub>6</sub>	Xaa <sub>7</sub>	Xaa <sub>8</sub>	Xaa <sub>9</sub>	Xaa <sub>10</sub>	Xaa <sub>11</sub>	Xaa <sub>12</sub>	Xaa <sub>13</sub>	Xaa <sub>14</sub>	Xaa <sub>15</sub>	Xaa <sub>16</sub>	Xaa <sub>17</sub>	Xaa <sub>18</sub>	Z
9	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
10	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
11	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
12	Tyr	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
13	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Tyr	NH <sub>2</sub>
14	His	Gly	Asp	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
15	His	Gly	Glu	naph	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
16	His	Gly	Glu	Phe	Ser	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
17	His	Gly	Glu	Phe	Ser	Thr	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
18	His	Gly	Glu	Phe	Thr	Thr	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
19	His	Gly	Glu	Phe	Thr	Ser	Glu	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
20	His	Gly	Glu	Phe	Thr	Ser	Asp	pGly	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
21	His	Gly	Glu	Phe	Thr	Ser	Asp	pGly	Leu	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
22	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	pGly	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>

FIGURE 10  
 (Sheet 1 of 2)

11/11

[SEQ. ID. NO.]	Xaa <sub>1</sub>	Xaa <sub>2</sub>	Xaa <sub>3</sub>	Xaa <sub>4</sub>	Xaa <sub>5</sub>	Xaa <sub>6</sub>	Xaa <sub>7</sub>	Xaa <sub>8</sub>	Xaa <sub>9</sub>	Xaa <sub>10</sub>	Xaa <sub>11</sub>	Xaa <sub>12</sub>	Xaa <sub>13</sub>	Xaa <sub>14</sub>	Xaa <sub>15</sub>	Xaa <sub>16</sub>	Xaa <sub>17</sub>	Xaa <sub>18</sub>	Z
23	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	pGly	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
24	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	naph	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
25	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Val	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
26	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Val	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH
27	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	tBuG	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
28	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	tBuG	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
29	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Asp	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
30	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
31	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	tPro	tPro	tPro	tPro	Ser	NH <sub>2</sub>
32	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	tPro	tPro	tPro	Ser	NH <sub>2</sub>
33	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	hPro	hPro	hPro	hPro	Ser	NH <sub>2</sub>
34	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	hPro	hPro	hPro	Ser	NH <sub>2</sub>
35	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	tPro	tPro	tPro	tPro	Ser	NH <sub>2</sub>
36	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	hPro	hPro	hPro	hPro	Ser	NH <sub>2</sub>
37	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	MeAla	MeAla	MeAla	MeAla	Ser	NH
38	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	MeAla	MeAla	MeAla	Ser	NH <sub>2</sub>
39	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	MeAla	MeAla	MeAla	MeAla	Ser	NH

FIGURE 10  
(Sheet 2 of 2)

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/00449

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 38/16

US CL : 514/2, 866

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/2, 866

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE, MEDLINE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,424,286 A (ENG) 13 June 1995 see entire document.	1-31

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

07 MAY 1998

Date of mailing of the international search report

29 MAY 1998

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Zohreh Fay

Telephone No. (703) 308-1235

**THIS PAGE BLANK (USPTO**



Effect of Exendin-4 and GLP-1 on Food Intake  
in NIH/SW Mice  
Mean  $\pm$  S.E.





FIGURE 1

**Effect of Exendin-4 on Food Intake in  
Female ob/ob Mice  
Mean  $\pm$  S.E.**



All injections given IP (5 ml/kg) at T=0 minutes

**FIGURE 2**







FIGURE 3

**Effect of Compound 1 on Food Intake  
in NIH/SW Mice  
Mean  $\pm$  S.E.**



**FIGURE 4**




FIGURE 5

Effect of Compound 3 on Food Intake  
In NIH/SW Mice  
Mean  $\pm$  S.E.




FIGURE 6

Effect of Compound 4 on Food Intake  
in NIH/SW Mice  
Mean  $\pm$  S.E.




FIGURE 7




FIGURE 8






FIGURE 9

1  
 2  
 3  
 4  
 5  
 6  
 7  
 8  
 9  
 10  
 11  
 12  
 13  
 14  
 15  
 16  
 17  
 18  
 19  
 20  
 21  
 22  
 23  
 24  
 25  
 26  
 27  
 28  
 29  
 30  
 31  
 32  
 33  
 34  
 35  
 36  
 37  
 38  
 39  
 40  
 41  
 42  
 43  
 44  
 45  
 46  
 47  
 48  
 49  
 50  
 51  
 52  
 53  
 54  
 55  
 56  
 57  
 58  
 59  
 60  
 61  
 62  
 63  
 64  
 65  
 66  
 67  
 68  
 69  
 70  
 71  
 72  
 73  
 74  
 75  
 76  
 77  
 78  
 79  
 80  
 81  
 82  
 83  
 84  
 85  
 86  
 87  
 88  
 89  
 90  
 91  
 92  
 93  
 94  
 95  
 96  
 97  
 98  
 99  
 100  
 101  
 102  
 103  
 104  
 105  
 106  
 107  
 108  
 109  
 110  
 111  
 112  
 113  
 114  
 115  
 116  
 117  
 118  
 119  
 120  
 121  
 122  
 123  
 124  
 125  
 126  
 127  
 128  
 129  
 130  
 131  
 132  
 133  
 134  
 135  
 136  
 137  
 138  
 139  
 140  
 141  
 142  
 143  
 144  
 145  
 146  
 147  
 148  
 149  
 150  
 151  
 152  
 153  
 154  
 155  
 156  
 157  
 158  
 159  
 160  
 161  
 162  
 163  
 164  
 165  
 166  
 167  
 168  
 169  
 170  
 171  
 172  
 173  
 174  
 175  
 176  
 177  
 178  
 179  
 180  
 181  
 182  
 183  
 184  
 185  
 186  
 187  
 188  
 189  
 190  
 191  
 192  
 193  
 194  
 195  
 196  
 197  
 198  
 199  
 200  
 201  
 202  
 203  
 204  
 205  
 206  
 207  
 208  
 209  
 210  
 211  
 212  
 213  
 214  
 215  
 216  
 217  
 218  
 219  
 220  
 221  
 222  
 223  
 224  
 225  
 226  
 227  
 228  
 229  
 230  
 231  
 232  
 233  
 234  
 235  
 236  
 237  
 238  
 239  
 240  
 241  
 242  
 243  
 244  
 245  
 246  
 247  
 248  
 249  
 250  
 251  
 252  
 253  
 254  
 255  
 256  
 257  
 258  
 259  
 260  
 261  
 262  
 263  
 264  
 265  
 266  
 267  
 268  
 269  
 270  
 271  
 272  
 273  
 274  
 275  
 276  
 277  
 278  
 279  
 280  
 281  
 282  
 283  
 284  
 285  
 286  
 287  
 288  
 289  
 290  
 291  
 292  
 293  
 294  
 295  
 296  
 297  
 298  
 299  
 300  
 301  
 302  
 303  
 304  
 305  
 306  
 307  
 308  
 309  
 310  
 311  
 312  
 313  
 314  
 315  
 316  
 317  
 318  
 319  
 320  
 321  
 322  
 323  
 324  
 325  
 326  
 327  
 328  
 329  
 330  
 331  
 332  
 333  
 334  
 335  
 336  
 337  
 338  
 339  
 340  
 341  
 342  
 343  
 344  
 345  
 346  
 347  
 348  
 349  
 350  
 351  
 352  
 353  
 354  
 355  
 356  
 357  
 358  
 359  
 360  
 361  
 362  
 363  
 364  
 365  
 366  
 367  
 368  
 369  
 370  
 371  
 372  
 373  
 374  
 375  
 376  
 377  
 378  
 379  
 380  
 381  
 382  
 383  
 384  
 385  
 386  
 387  
 388  
 389  
 390  
 391  
 392  
 393  
 394  
 395  
 396  
 397  
 398  
 399  
 400  
 401  
 402  
 403  
 404  
 405  
 406  
 407  
 408  
 409  
 410  
 411  
 412  
 413  
 414  
 415  
 416  
 417  
 418  
 419  
 420  
 421  
 422  
 423  
 424  
 425  
 426  
 427  
 428  
 429  
 430  
 431  
 432  
 433  
 434  
 435  
 436  
 437  
 438  
 439  
 440  
 441  
 442  
 443  
 444  
 445  
 446  
 447  
 448  
 449  
 450  
 451  
 452  
 453  
 454  
 455  
 456  
 457  
 458  
 459  
 460  
 461  
 462  
 463  
 464  
 465  
 466  
 467  
 468  
 469  
 470  
 471  
 472  
 473  
 474  
 475  
 476  
 477  
 478  
 479  
 480  
 481  
 482  
 483  
 484  
 485  
 486  
 487  
 488  
 489  
 490  
 491  
 492  
 493  
 494  
 495  
 496  
 497  
 498  
 499  
 500  
 501  
 502  
 503  
 504  
 505  
 506  
 507  
 508  
 509  
 510  
 511  
 512  
 513  
 514  
 515  
 516  
 517  
 518  
 519  
 520  
 521  
 522  
 523  
 524  
 525  
 526  
 527  
 528  
 529  
 530  
 531  
 532  
 533  
 534  
 535  
 536  
 537  
 538  
 539  
 540  
 541  
 542  
 543  
 544  
 545  
 546  
 547  
 548  
 549  
 550  
 551  
 552  
 553  
 554  
 555  
 556  
 557  
 558  
 559  
 560  
 561  
 562  
 563  
 564  
 565  
 566  
 567  
 568  
 569  
 570  
 571  
 572  
 573  
 574  
 575  
 576  
 577  
 578  
 579  
 580  
 581  
 582  
 583  
 584  
 585  
 586  
 587  
 588  
 589  
 590  
 591  
 592  
 593  
 594  
 595  
 596  
 597  
 598  
 599  
 600  
 601  
 602  
 603  
 604  
 605  
 606  
 607  
 608  
 609  
 610  
 611  
 612  
 613  
 614  
 615  
 616  
 617  
 618  
 619  
 620  
 621  
 622  
 623  
 624  
 625  
 626  
 627  
 628  
 629  
 630  
 631  
 632  
 633  
 634  
 635  
 636  
 637  
 638  
 639  
 640  
 641  
 642  
 643  
 644  
 645  
 646  
 647  
 648  
 649  
 650  
 651  
 652  
 653  
 654  
 655  
 656  
 657  
 658  
 659  
 660  
 661  
 662  
 663  
 664  
 665  
 666  
 667  
 668  
 669  
 670  
 671  
 672  
 673  
 674  
 675  
 676  
 677  
 678  
 679  
 680  
 681  
 682  
 683  
 684  
 685  
 686  
 687  
 688  
 689  
 690  
 691  
 692  
 693  
 694  
 695  
 696  
 697  
 698  
 699  
 700  
 701  
 702  
 703  
 704  
 705  
 706  
 707  
 708  
 709  
 710  
 711  
 712  
 713  
 714  
 715  
 716  
 717  
 718  
 719  
 720  
 721  
 722  
 723  
 724  
 725  
 726  
 727  
 728  
 729  
 730  
 731  
 732  
 733  
 734  
 735  
 736  
 737  
 738  
 739  
 740  
 741  
 742  
 743  
 744  
 745  
 746  
 747  
 748  
 749  
 750  
 751  
 752  
 753  
 754  
 755  
 756  
 757  
 758  
 759  
 760  
 761  
 762  
 763  
 764  
 765  
 766  
 767  
 768  
 769  
 770  
 771  
 772  
 773  
 774  
 775  
 776  
 777  
 778  
 779  
 780  
 781  
 782  
 783  
 784  
 785  
 786  
 787  
 788  
 789  
 790  
 791  
 792  
 793  
 794  
 795  
 796  
 797  
 798  
 799  
 800  
 801  
 802  
 803  
 804  
 805  
 806  
 807  
 808  
 809  
 810  
 811  
 812  
 813  
 814  
 815  
 816  
 817  
 818  
 819  
 820  
 821  
 822  
 823  
 824  
 825  
 826  
 827  
 828  
 829  
 830  
 831  
 832  
 833  
 834  
 835  
 836  
 837  
 838  
 839  
 840  
 841  
 842  
 843  
 844  
 845  
 846  
 847  
 848  
 849  
 850  
 851  
 852  
 853  
 854  
 855  
 856  
 857  
 858  
 859  
 860  
 861  
 862  
 863  
 864  
 865  
 866  
 867  
 868  
 869  
 870  
 871  
 872  
 873  
 874  
 875  
 876  
 877  
 878  
 879  
 880  
 881  
 882  
 883  
 884  
 885  
 886  
 887  
 888  
 889  
 890  
 891  
 892  
 893  
 894  
 895  
 896  
 897  
 898  
 899  
 900  
 901  
 902  
 903  
 904  
 905  
 906  
 907  
 908  
 909  
 910  
 911  
 912  
 913  
 914  
 915  
 916  
 917  
 918  
 919  
 920  
 921  
 922  
 923  
 924  
 925  
 926  
 927  
 928  
 929  
 930  
 931  
 932  
 933  
 934  
 935  
 936  
 937  
 938  
 939  
 940  
 941  
 942  
 943  
 944  
 945  
 946  
 947  
 948  
 949  
 950  
 951  
 952  
 953  
 954  
 955  
 956  
 957  
 958  
 959  
 960  
 961  
 962  
 963  
 964  
 965  
 966  
 967  
 968  
 969  
 970  
 971  
 972  
 973  
 974  
 975  
 976  
 977  
 978  
 979  
 980  
 981  
 982  
 983  
 984  
 985  
 986  
 987  
 988  
 989  
 990  
 991  
 992  
 993  
 994  
 995  
 996  
 997  
 998  
 999  
 1000  
 1001  
 1002  
 1003  
 1004  
 1005  
 1006  
 1007  
 1008  
 1009  
 1010  
 1011  
 1012  
 1013  
 1014  
 1015  
 1016  
 1017  
 1018  
 1019  
 1020  
 1021  
 1022  
 1023  
 1024  
 1025  
 1026  
 1027  
 1028  
 1029  
 1030  
 1031  
 1032  
 1033  
 1034  
 1035  
 1036  
 1037  
 1038  
 1039  
 1040  
 1041  
 1042  
 1043  
 1044  
 1045  
 1046  
 1047  
 1048  
 1049  
 1050  
 1051  
 1052  
 1053  
 1054  
 1055  
 1056  
 1057  
 1058  
 1059  
 1060  
 1061  
 1062  
 1063  
 1064  
 1065  
 1066  
 1067  
 1068  
 1069  
 1070  
 1071  
 1072  
 1073  
 1074  
 1075  
 1076  
 1077  
 1078  
 1079  
 1080  
 1081  
 1082  
 1083  
 1084  
 1085  
 1086  
 1087  
 1088  
 1089  
 1090  
 1091  
 1092  
 1093  
 1094  
 1095  
 1096  
 1097  
 1098  
 1099  
 1100  
 1101  
 1102  
 1103  
 1104  
 1105  
 1106  
 1107  
 1108  
 1109  
 1110  
 1111  
 1112  
 1113  
 1114  
 1115  
 1116  
 1117  
 1118  
 1119  
 1120  
 1121  
 1122  
 1123  
 1124  
 1125  
 1126  
 1127  
 1128  
 1129  
 1130  
 1131  
 1132  
 1133  
 1134  
 1135  
 1136  
 1137  
 1138  
 1139  
 1140  
 1141  
 1142  
 1143  
 1144  
 1145  
 1146  
 1147  
 1148  
 1149  
 1150  
 1151  
 1152  
 1153  
 1154  
 1155  
 1156  
 1157  
 1158  
 1159  
 1160  
 1161  
 1162  
 1163  
 1164  
 1165  
 1166  
 1167  
 1168  
 1169  
 1170  
 1171  
 1172  
 1173  
 1174  
 1175  
 1176  
 1177  
 1178  
 1179  
 1180  
 1181  
 1182  
 1183  
 1184  
 1185  
 1186  
 1187  
 1188  
 1189  
 1190  
 1191  
 1192  
 1193  
 1194  
 1195  
 1196  
 1197  
 1198  
 1199  
 1200  
 1201  
 1202  
 1203  
 1204  
 1205  
 1206  
 1207  
 1208  
 1209  
 1210  
 1211  
 1212  
 1213  
 1214  
 1215  
 1216  
 1217  
 1218  
 1219  
 1220  
 1221  
 1222  
 1223  
 1224  
 1225  
 1226  
 1227  
 1228  
 1229  
 1230  
 1231  
 1232  
 1233  
 1234  
 1235  
 1236  
 1237  
 1238  
 1239  
 1240  
 1241  
 1242  
 1243  
 1244  
 1245  
 1246  
 1247  
 1248  
 1249  
 1250  
 1251  
 1252  
 1253  
 1254  
 1255  
 1256  
 1257  
 1258  
 1259  
 1260  
 1261  
 1262  
 1263  
 1264  
 1265  
 1266  
 1267  
 1268  
 1269  
 1270  
 1271  
 1272  
 1273  
 1274  
 1275  
 1276  
 1277  
 1278  
 1279  
 1280  
 1281  
 1282  
 1283  
 1284  
 1285  
 1286  
 1287  
 1288  
 1289  
 1290  
 1291  
 1292  
 1293  
 1294  
 1295  
 1296  
 1297  
 1298  
 1299  
 1300  
 1301  
 1302  
 1303  
 1304  
 1305  
 1306  
 1307  
 1308  
 1309  
 1310  
 1311  
 1312  
 1313  
 1314  
 1315  
 1316  
 1317  
 1318  
 1319  
 1320  
 1321  
 1322  
 1323  
 1324  
 1325  
 1326  
 1327  
 1328  
 1329  
 1330  
 1331  
 1332  
 1333  
 1334  
 1335  
 1336  
 1337  
 1338  
 1339  
 1340  
 1341  
 1342  
 1343  
 1344  
 1345  
 1346  
 1347  
 1348  
 1349  
 1350  
 1351  
 1352  
 1353  
 1354  
 1355  
 1356  
 1357  
 1358  
 1359  
 1360  
 1361  
 1362  
 1363  
 1364  
 1365  
 1366  
 1367  
 1368  
 1369  
 1370  
 1371  
 1372  
 1373  
 1374  
 1375  
 1376  
 1377  
 1378  
 1379  
 1380  
 1381  
 1382  
 1383  
 1384  
 1385  
 1386  
 1387  
 1388  
 1389  
 1390  
 1391  
 1392  
 1393  
 1394  
 1395  
 1396  
 1397  
 1398  
 1399  
 1400  
 1401  
 1402  
 1403  
 1404  
 1405  
 1406  
 1407  
 1408  
 1409  
 1410  
 1411  
 1412  
 1413  
 1414  
 1415  
 1416  
 1417  
 1418  
 1419  
 1420  
 1421  
 1422  
 1423  
 1424  
 1425  
 1426  
 1427  
 1428  
 1429  
 1430  
 1431  
 1432  
 1433  
 1434  
 1435  
 1436  
 1437  
 1438  
 1439  
 1440  
 1441  
 1442  
 1443  
 1444  
 1445  
 1446  
 1447  
 1448  
 1449  
 1450  
 1451  
 1452  
 1453  
 1454  
 1455  
 1456  
 1457  
 1458  
 1459  
 1460  
 1461  
 1462  
 1463  
 1464  
 1465  
 1466  
 1467  
 1468  
 1469  
 1470  
 1471  
 1472  
 1473  
 1474  
 1475  
 1476  
 1477  
 1478  
 1479  
 1480  
 1481  
 1482  
 1483  
 1484  
 1485  
 1486  
 1487



**THIS PAGE BLANK (USPTO)**